Additional Perspectives articles for *Influenza: The Cutting Edge* book collection are available at http://perspectivesinmedicine.cshlp.org/cgi/collection/influenza_the_cutting_edge.

Structure and Function of the Influenza Virus Transcription and Replication Machinery

Ervin Fodor¹ and Aartjan J.W. te Velthuis²

¹Sir William Dunn School of Pathology, University of Oxford, Oxford OX1 3RE, United Kingdom

²Division of Virology, Department of Pathology, University of Cambridge, Cambridge CB2 2QQ, United Kingdom

Correspondence: ervin.fodor@path.ox.ac.uk; ajwt6@cam.ac.uk

Transcription and replication of the influenza virus RNA genome is catalyzed by the viral heterotrimeric RNA-dependent RNA polymerase in the context of viral ribonucleoprotein (vRNP) complexes. Atomic resolution structures of the viral RNA synthesis machinery have offered insights into the initiation mechanisms of viral transcription and genome replication, and the interaction of the viral RNA polymerase with host RNA polymerase II, which is required for the initiation of viral transcription. Replication of the viral RNA genome by the viral RNA polymerase depends on host ANP32A, and host-specific sequence differences in ANP32A underlie the poor activity of avian influenza virus polymerases in mammalian cells. A failure to faithfully copy the viral genome segments can lead to the production of aberrant viral RNA products, such as defective interfering (DI) RNAs and mini viral RNAs (mvRNAs). Both aberrant RNA types have been implicated in innate immune responses against influenza virus infection. This review discusses recent insights into the structure–function relationship of the viral RNA polymerase and its role in determining host range and virulence.

Infections with influenza viruses type A and B are responsible for annual seasonal epidemics leading to about half a million deaths worldwide. Influenza type A viruses also cause occasional pandemics, which can claim millions of lives. In contrast, influenza C type infections are milder and influenza type D viruses, which infect pigs and cattle, have not been associated with human disease (Krammer et al. 2018; Taubenberger et al. 2019).

The influenza virus genome consists of eight (influenza A and B viruses) or seven (influenza C and D viruses) segments of single-stranded negative-sense viral RNA (vRNA) (Fig. 1A).

Each segment consists of conserved, partially complementary 5' and 3' termini, segment-specific untranslated regions (UTRs) and one or more open-reading frames (ORFs) (Fig. 1A). The vRNA segments are assembled into viral ribonucleoprotein (vRNP) complexes, in which the 5' and 3' termini are bound by a viral RNA-dependent RNA polymerase (RdRp) and the rest of the RNA is bound by oligomeric viral nucleoprotein (NP), forming a double-helical rod-like structure (Fig. 1B,C; Eisfeld et al. 2015). Internal base-pairing within the segments allows the formation of secondary RNA structures (Ferhadian et al. 2018). Influenza viruses must express a full

Editors: Gabriele Neumann and Yoshihiro Kawaoka

Additional Perspectives on Influenza: The Cutting Edge available at www.perspectivesinmedicine.org

Copyright © 2020 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a038398

Cite this article as *Cold Spring Harb Perspect Med* 2020;10:a038398

E. Fodor and A.J.W. te Velthuis

Figure 1. Influenza virus RNAs and viral ribonucleoprotein (vRNP) structure. (A) Influenza virus viral RNAs (vRNAs). The influenza A virus genome consists of eight segments of single-stranded negative sense vRNA. Each segment contains conserved 5' and 3' termini, segment-specific untranslated regions (UTRs), and at least one large open-reading frame (ORF). (B) Schematic of influenza virus vRNP and transcription and replication. vRNA segments are assembled into vRNPs with multiple copies of nucleoprotein (NP) and a single copy of the viral RNA-dependent RNA polymerase (RdRp), composed of PB1, PB2, and PA subunits, that binds the conserved 5' and 3' vRNA termini. The RNA polymerase, in the context of vRNPs, transcribes vRNA into viral mRNA, which contains a 5' cap-1 structure (cap) and a 3' poly(A) tail (A_n), and replicates vRNA through a complementary RNA (cRNA) replicative intermediate, which is assembled with NP and polymerase into complementary ribonucleoprotein (cRNP). (C) Structure of the influenza virus vRNP. NP protomers are shown in surface and cartoon representation (PDB 4BBL).

complement of their genome segments in the infected cell to be productively infectious. The segments can be contributed by one virion or, in the case of a defective particle, multiple virions.

After fusion of viral and endosomal membranes during influenza virus entry, vRNPs are released into the cytoplasm and transported into the nucleus in which viral transcription takes place (Walker and Fodor 2019). It is in the context of nuclear vRNPs that the viral RNA polymerase transcribes the vRNA into positive-sense

capped and polyadenylated mRNA for translation of viral proteins (Fig. 1B). The influenza A virus genome encodes 10 major and several auxiliary proteins. Viral protein expression is essential for the replication of the viral RNA genome by the viral RNA polymerase, which proceeds through a complementary RNA (cRNA) replicative intermediate (Fig. 1B; te Velthuis and Fodor 2016; Pflug et al. 2017).

Recent structural and functional studies of the influenza RNA synthesis machinery have

provided novel insights into the mechanisms by which influenza viruses transcribe and replicate their RNA genome (te Velthuis and Fodor 2016; Pflug et al. 2017; Lo et al. 2018). These studies have answered fundamental questions about influenza RNA polymerase function and contributed to our understanding of the dependence of the influenza virus on host factors for the transcription and replication of its genome. This review discusses how RNA polymerase structures inform us about the function of the RNA synthesis machinery and its role in defining viral virulence and host range.

ARCHITECTURE OF vRNPs

vRNPs are composed of one single-stranded vRNA segment, a copy of the RNA polymerase bound to the partially complementary 5' and 3' vRNA termini, and multiple copies of NP that are arranged in a double helical conformation (Fig. 1B,C). Currently, only low-resolution structural models of vRNPs are available that differ in their helical parameters and handedness and do not allow unambiguous mapping of the path of the RNA and NP–NP contacts within vRNPs (Arranz et al. 2012; Moeller et al. 2012). Studies aimed at obtaining high-resolution vRNP structures are hampered by the heterogeneity in vRNA length (890–2341 nucleotides for influenza A virus; Fig. 1A) and high-plasticity of the NP–NP interactions, which leads to the formation of curves and kinks in vRNPs, and complicates computational processing of cryogenic electron microscopy (cryo-EM) data (Gallagher et al. 2017).

NP is a crescent-shaped molecule that is comprised of a head and body domain and a 28 amino acid–long tail loop. NP forms oligomers when the tail loop of one NP inserts into the body domain of a neighboring NP (Ye et al. 2006; Ng et al. 2008; Turrell et al. 2013). A basic groove separating the head and body domains represents the RNA-binding surface. Phosphorylation and dephosphorylation of NP by cellular kinases and phosphatases affects NP oligomerization and viral replication (Hutchinson et al. 2012; Chenavas et al. 2013; Mondal et al. 2015, 2017; Turrell et al. 2015). The footprint of NP on

vRNA is ~24 nucleotides (Ortega et al. 2000; Hutchinson et al. 2012), but it is not known how many nucleotides make direct contact with NP and how the RNA is arranged in vRNPs. Ribonuclease digestion studies show that the RNA in vRNP is susceptible to hydrolytic cleavage suggesting that vRNA is bound on the outside surface of NP. Mapping studies of NP–vRNA interactions have identified regions of low and high NP density (Lee et al. 2017; Williams et al. 2018), and chemical probing of the vRNA structure in vRNPs shows that vRNPs packaged into virions can accommodate secondary structures in their vRNAs (Dadonaite et al. 2019), showing that NP does not associate with vRNA evenly. The NP helix in vRNPs thus likely provides a scaffold for the formation of secondary RNA structures. These RNA structures can facilitate intersegments RNA–RNA interactions and selective packaging of the vRNPs into virions (Dadonaite et al. 2019).

ARCHITECTURE OF THE RNA POLYMERASE

The influenza virus RNA polymerase is composed of the proteins polymerase basic 1 (PB1), PB2, and polymerase acidic (PA) (P3 in influenza C and D viruses) (Fig. 2A; te Velthuis and Fodor 2016; Pflug et al. 2017; Lo et al. 2018). X-ray crystallography and cryo-EM studies showed that the RNA polymerase consists of a core domain and several peripheral domains that are connected to the core through highly flexible linkers (Pflug et al. 2014; Reich et al. 2014; Chang et al. 2015; Hengrung et al. 2015; Thierry et al. 2016; Fan et al. 2019; Peng et al. 2019). The PB1 subunit contributes most to the core domain, and contains the fingers, palm, and thumb subdomains that are characteristic of RNA polymerases, and a priming loop, a β -hairpin structure that protrudes from the thumb domain into the polymerase active site and that is typical of RNA polymerases that can initiate RNA synthesis de novo (Fig. 2B; te Velthuis 2014). The PA subunit is composed of two domains. The amino-terminal domain contains endonuclease activity and is connected to the polymerase core via a flexible linker. In contrast, the carboxy-terminal domain is intimately

E. Fodor and A.J.W. te Velthuis

Figure 2. Structure of the influenza virus RNA polymerase. (A) Surface presentation of the influenza A virus RNA polymerase (PDB 6RR7) bound to capped RNA primer and 5' and 3' vRNA termini. The subunits of the RNA polymerase are colored as in Figure 1. (B) Schematic model of the influenza A virus RNA polymerase showing the core chamber of the polymerase and its connection to the solvent via template and nucleotide triphosphate (NTP) entry channels, and template and product exit channels. The location of the priming loop (PL) as well as the 3' and 5' end-binding pockets are indicated.

associated with PB1 and tightly integrated into the thumb subdomain of the polymerase core. The amino-terminal third of PB2 is also part of the polymerase core, but the carboxy-terminal two thirds form multiple flexible domains, including the cap-binding, midlink, 627, and nuclear localization signal (NLS) domains. All subunits contribute to the formation of the channels that facilitate template RNA entry and exit, nucleotide triphosphate (NTP) entry, and product RNA exit (Fig. 2B).

The polymerase contains several sites for binding viral RNA (Fig. 2). The 5' end of vRNA and cRNA are tightly bound near the template entry channel of the polymerase by a pocket composed of amino acid residues contributed by the PB1 and PA subunits (Pflug et al. 2014; Reich et al. 2014; Robb et al. 2016; Fan et al. 2019; Peng et al. 2019). The 3' end of the vRNA can associate with two different sites (the A-site and B-site) on the surface of the polymerase, whereas the 3' end of cRNA has been shown to associate with the B-site only (Fig. 2B; Pflug et al. 2014; Reich et al. 2014; Fan et al. 2019; Peng et al. 2019). The functional significance of these surface 3'-end-binding sites remains to be determined, but they likely provide (1) a preinitiation site from which the 3' end of vRNA enters the polymerase active site (the A-site), (2) a preinitiation site from which the 3' end of cRNA enters the polymerase active site (the B-site), or (3) a site to accommodate the 3' end of the RNA

template once it emerges from the polymerase active site after being copied (the B-site).

TRANSCRIPTION

Transcription is a primer-dependent process that involves the generation of 5' capped and 3' polyadenylated mRNA molecules using the negative-sense vRNA segments as templates by the viral RNA polymerase (Fig. 1B). Unlike the RNA polymerases from nonsegmented negative-sense RNA viruses, the influenza virus RNA polymerase cannot synthesize and methylate cap structures. To ensure that influenza virus mRNAs contain a cap-1 structure at their 5' terminus, the RNA polymerase needs to perform several key steps. First, the RNA polymerase associated with vRNPs, that is, the resident polymerase, binds the carboxy-terminal domain (CTD) of the large subunit of host RNA polymerase II (Pol II) to bring vRNPs in close proximity to nascent host 5' capped RNAs (Fig. 3A; Engelhardt et al. 2005; Martínez-Alonso et al. 2016; Lukarska et al. 2017; Serna Martin et al. 2018). Pol II CTD interacts with the viral RNA polymerase at several binding sites (Lukarska et al. 2017; Serna Martin et al. 2018), which stabilizes the PB2 cap-binding and PA endonuclease domains in a transcription-ready conformation (Serna Martin et al. 2018). Second, the PB2 cap-binding domain binds the 5' cap of the nascent host RNA and the PA endonuclease

A

Pol II

CTD

cap

DNA

PB2 cap

PA endo

B

Preinitiation

5'

3'

PL

PB2 cap

Elongation

5'

9-bp duplex

Duplex melting

PB2 cap

domain cleaves the mRNA 10–14 bases downstream from the 5' cap (Fig. 3A). Third, the 3' end of the resulting capped RNA primer is inserted into the polymerase active site where it base-pairs with the 3' end of the vRNA template (Fig. 3B). Initiation of viral mRNA synthesis occurs at the penultimate or third base from the 3' end of the vRNA, depending on where the 3' end of the primer has base-paired with the vRNA (Pflug et al. 2018; te Velthuis and Oymans 2018; Kouba et al. 2019). A poor transition from initiation to elongation may result in early termination at the fourth base from the 3' end of the vRNA. The RNA polymerase can use the

Cite this article as *Cold Spring Harb Perspect Med* 2020;10:a038398

E. Fodor and A.J.W. te Velthuis

product are directed toward their respective exit channels (Kouba et al. 2019). Finally, termination of the viral transcription product occurs by stuttering of the RNA polymerase on a sequence of uridine residues near the 5' end of the vRNA template, leading to the addition of a poly(A) tail (Robertson et al. 1981; Poon et al. 1999).

REPLICATION

Replication of the vRNA segments by the viral RNA polymerase is a two-step process (Fig. 4A, B). In the first step, vRNA is copied into a cRNA replicative intermediate (Fig. 4A). This step is initiated by entry of the 3' end into the active

Figure 4. Replication by the influenza virus RNA polymerase. (A) Synthesis of complementary RNA (cRNA) by the influenza virus RNA polymerase and coreplicative assembly of complementary ribonucleoprotein (cRNP). (B) Synthesis of viral RNA (vRNA) by the influenza virus RNA polymerase and coreplicative assembly of vRNP. Template realignment is promoted by a regulatory polymerase. ANP32A associates with RNA-free RNA polymerase and may regulate dimerization of the RNA polymerase. (C) Schematic model of the influenza virus RNA polymerase showing replication initiation during cRNA synthesis. (D) Schematic model of the influenza virus RNA polymerase showing replication initiation and realignment during vRNA synthesis.

site, presumably from the A-site (Fig. 2B; Robb et al. 2019). Next, the viral polymerase uses de novo initiation at the 3' terminal UC of the vRNA template to generate a pppApG dinucleotide (Fig. 4C). Initiation is dependent on the priming loop, which contributes to the stabilization of the initiation complex (te Velthuis et al. 2016). After initiation, the priming loop is likely extruded from the active site to make space for extension of the nascent cRNA, as has been observed during transcription elongation (Kouba et al. 2019), because truncation of the priming loop stimulates elongation (te Velthuis et al. 2016). As the nascent cRNA emerges from the RNA polymerase, the cRNA assembles into a complementary ribonucleoprotein (cRNP) complex, a vRNP-like structure, in which a newly produced RNA polymerase associates with the terminal cRNA ends and oligomeric NP binds the rest of the cRNA stabilizing it from degradation (Fig. 4A; Vreede et al. 2004; York et al. 2013). Expression of RNA polymerase and NP is indeed essential for the accumulation of cRNA during infection.

In the second step of replication, cRNA in the context of cRNPs acts as a template for the synthesis of vRNA (Fig. 4B). The 3' end of the cRNA has been shown to bind at the B-site on the surface of the polymerase (Fig. 2B), which suggests that the 3' cRNA terminus uses a different preinitiation site before the 3' end of the cRNA enters the active site and initiation occurs (Fan et al. 2019; Peng et al. 2019). In contrast to initiation on the vRNA template, the polymerase initiates internally at positions 4 and 5 of the cRNA template to produce a pppApG dinucleotide (Fig. 4D). The dinucleotide next acts as a primer for terminal initiation at positions 1 and 2 of cRNA after backtracking of the cRNA template (Deng et al. 2006) with the aid of the priming loop and a regulatory polymerase (Oymans and te Velthuis 2018; Fan et al. 2019). The regulatory polymerase forms a dimer with the resident cRNP-associated polymerase, which carries out replication. Several studies have shown that the influenza virus RNA polymerase can dimerize and that genome replication requires a second polymerase in addition to the polymerase that resides in the cRNP (Jorba

et al. 2008; York et al. 2013; Chang et al. 2015; Fan et al. 2019; Peng et al. 2019). In the influenza A virus polymerase, PA residues 352–356 of both polymerases, and PB2 residues 71–76 of one polymerase and PA residue 347 of the second polymerase contribute to the dimer interface (Chang et al. 2015; Fan et al. 2019). Mutation of the dimer interface or its disruption with a nanobody inhibit vRNA synthesis by reducing the efficiency of cRNA template backtracking, and thereby realignment of the pppApG and terminal pppApG-primed initiation (Fan et al. 2019). Specifically, analysis of the priming loop shows that the β -hairpin adopts different positions dependent on the oligomeric state of the polymerase. When the polymerase is in a monomeric form, the priming loop resides in a position that would promote internal initiation, but on dimerization the priming loop changes position and facilitates backtracking of the template RNA to allow terminal pppApG-primed initiation of vRNA synthesis (Fan et al. 2019). The 5' end of the vRNA product is most likely captured by a stabilizing polymerase as soon as it emerges from the active site of the replicating cRNP-resident polymerase, initiating the assembly of the vRNA with NP into vRNP (Fig. 4B). Whether it is the same or two different polymerase heterotrimers that carry out regulation and stabilization remains to be determined.

HOST FACTORS AND ADAPTIVE MUTATIONS

Influenza virus polymerase function depends on association with host factors (Peacock et al. 2019). The transcriptional activity of the viral RNA polymerase is dependent on host Pol II for capped RNA primers that facilitate transcription initiation (Plotch et al. 1981) and structural studies have shown that the interaction of the viral polymerase with the CTD of Pol II is important for stabilizing a conformation of the viral polymerase that is competent in cap-snatching and transcription initiation (Fig. 3A; Lukarska et al. 2017; Serna Martin et al. 2018). ANP32 proteins have been identified as further host factors that are essential for viral polymerase activity. Simultaneous knockout of both

E. Fodor and A.J.W. te Velthuis

ANP32A and ANP32B isoforms in human cells results in cells that are unable to support influenza virus polymerase activity in minigenome assays and virus growth (Staller et al. 2019; Zhang et al. 2019). ANP32 proteins have been linked to viral genome replication, particularly the second step involving the synthesis of vRNA from the cRNA template (Fig. 4B; Sugiyama et al. 2015), suggesting that ANP32 proteins are important factors for the replicase function of the polymerase. ANP32A can directly interact with the apo form of the viral polymerase (Sugiyama et al. 2015; Baker et al. 2018; Staller et al. 2019), in agreement with a hypothesis that ANP32A may help recruit a second polymerase for vRNP formation. However, the molecular details of how ANP32A protein interacts with the replicase remain unknown.

The dependence of the viral polymerase on ANP32 proteins plays a critical role in the host restriction of avian influenza virus polymerases in mammalian cells (Long et al. 2016). A key determinant of host range is PB2 amino acid 627. This residue is a glutamic acid in avian influenza virus polymerases (Subbarao et al. 1993), but an adaptive mutation to lysine allows the avian influenza polymerase to function efficiently in a mammalian cell. In addition to the E627K change, alternative adaptive mutations have been observed. Most notably, the swine origin 2009 pandemic virus retained 627E in its avian PB2 gene and used an adaptation at position 591 to increase viral replication (Mehle and Doudna 2009; Yamada et al. 2010). These adaptive changes are linked to a host-specific difference in the ANP32 proteins. Specifically, avian ANP32A contains a 33-amino-acids insertion downstream from amino acid 175. This insertion, which consists of six unique amino acids representing a hydrophobic SUMO interaction motif (SIM)-like sequence and a 27-amino-acid-long duplication of amino acids 149 to 175, is critical for the support of PB2 627E-containing polymerase (Long et al. 2016; Domingues and Hale 2017; Baker et al. 2018). Human ANP32A and ANP32B variants as well avian ANP32A splice forms that lack this sequence are unable to support the activity of an avian-adapted influenza polymerase with PB2 627E. Currently,

the molecular basis of how the E627K mutation allows the polymerase to work with the mammalian version of ANP32A proteins remains unknown.

DI RNAs, mvRNAs, AND svRNAs

In addition to the synthesis of full-length copies of the vRNA segments, influenza virus replication produces defective interfering (DI) RNAs, mini viral RNAs (mvRNAs) and small viral RNAs (svRNAs) (Fig. 5A). DI RNAs were first described in the context of DI particles (DIPs), which are virions that are able to interfere with the production of wild-type virus (Dimmock and Easton 2014). The DI RNAs have large internal deletions but usually retain at least 150–200 nucleotides from both termini, including the conserved 5' and 3' terminal sequences (Jennings et al. 1983). DI RNAs are encapsidated by multiple NP molecules and are bound by a single copy of the RNA polymerase, similar to full-length viral genome segments (Fig. 5A; Coloma et al. 2009). DI RNAs retain sufficient sequence information to be packaged into virions replacing the parent full-length vRNA segment.

Next-generation sequencing of DI RNAs has shown that deletions can occur in any genome segment, but that some segments are more prone to give rise to DI RNAs than other segments. Specifically, DI RNAs derived from the three polymerase encoding segments are most frequently observed (Saira et al. 2013; Al-naji et al. 2019). DI RNAs are generated through a still poorly defined mechanism, in which the viral polymerase ceases product elongation at one site of the viral RNA template, only to resume elongation at a downstream site, which results in a failure to copy the interlaying stretch of the template (Fig. 5B). It has been hypothesized that an intramolecular copy-choice mechanism could be involved, in which resumption of RNA synthesis could be dependent on base-pairing between the 3' end of the nascent product and the RNA template. However, a recent analysis of the sequences that flank the deletion (i.e., polymerase stop and restart site) found no evidence for the enrichment of direct repeat sequences at junctions, which argues against a significant role

Influenza Virus RNA Polymerase Function

Figure 5. Aberrant RNA synthesis by the influenza virus RNA polymerase. (A) In addition to full-length viral RNAs (vRNAs), the influenza virus RNA polymerase produces defective interfering (DI) RNAs, mini viral RNAs (mvRNAs), and small viral RNAs (svRNAs). (B) Synthesis of DI RNAs and mvRNAs uses an intramolecular copy-choice mechanism. (C) Host-specific mutation in the viral RNA polymerase are linked to mvRNA levels. mvRNAs can be bound by RIG-I and induce RIG-I activation and cytokine expression.

for direct repeats in DI RNA formation (Alnaji et al. 2019). Interestingly, DI RNAs encode open reading frames and could express viral protein fragments or novel proteins that may have a function in the viral replication cycle (Akkina et al. 1984; Boergeling et al. 2015).

mvRNAs are ~50–125-nt-long aberrant replication products with very large internal deletions, retaining the conserved 5' and 3' termini (Fig. 5A). In contrast to DI RNAs, mvRNAs can be replicated and transcribed in the absence of NP (Turell et al. 2013; te Velhuis et al. 2018). Consequently, they are not (fully) encapsidated

and likely not packaged into virions. Formation of mvRNAs appears to follow an intramolecular copy-choice mechanism that does not require perfect complementarity similar to DI formation (Fig. 5B) and that is more prevalent during the replication of some vRNA segments than others, suggesting a dependence on RNA sequence and/or structure (te Velhuis et al. 2018). The generation of mvRNAs has been linked to the dysregulation of replication, for example by an imbalance in the polymerase–NP ratio, as well as to specific amino acids in the polymerase (te Velhuis et al. 2018; Liu

E. Fodor and A.J.W. te Velthuis

et al. 2019). Specifically, an amino acid change increasing polymerase fidelity has been shown to decrease mvRNA production, whereas amino acids characteristic of avian influenza A virus polymerases increase mvRNA production in mammalian cells (te Velthuis et al. 2018).

In contrast to DI RNAs and mvRNAs, svRNAs are 22–27 nt long RNAs that only contain the 5' terminus of vRNAs (Fig. 5A). They likely represent early termination products of vRNA synthesis from a cRNA template. svRNAs have been found in cultured infected cells as well as in infected mouse lung cells, and may play a role in regulating viral replication (Perez et al. 2010, 2012; Koiré et al. 2016). In particular, it has been shown that svRNA synthesis can influence vRNA synthesis by binding to the viral RNA polymerase (Perez et al. 2012). The viral nonstructural protein 2 (NS2/NEP) may be involved in regulating svRNA expression (Perez et al. 2010; Liu et al. 2019).

INNATE IMMUNE RESPONSES

The innate immune response plays a critical role in the outcome of influenza A virus infections. In particular, a dysregulated immune response underpins the high-virulence of pandemic and avian influenza virus infections in humans (de Jong et al. 2006; Kash et al. 2006; Forero et al. 2016). Key cytokines involved in influenza A virus infections include type I and III interferons, such as interferon β and λ (Klinkhammer et al. 2018). These cytokines are typically expressed in response to the detection of viral RNA by the cellular pathogen sensor retinoic acid-inducible gene I (RIG-I), which is activated by binding short double-stranded RNA containing a 5' di- or triphosphate. Influenza vRNA and cRNA contain 5' triphosphate and partially complementary termini, which could be bound by RIG-I (Yoneyama et al. 2004; Kato et al. 2006; Pichlmair et al. 2006). In agreement with viral RNA being important for RIG-I activation, correlation between the activity of polymerase and the expression of cytokines has been observed in a number of studies (Forero et al. 2016; Li et al. 2018; Russell et al. 2019). Specifically, the polymerases of the 1918 H1N1 pandemic virus and

H5N1 avian influenza viruses have been linked to excessive cytokine production (Forero et al. 2016; Li et al. 2018).

Genomic viral RNA has been shown to be a poor inducer of RIG-I activation (Rehwinkel et al. 2010), likely because the binding of the viral RNA polymerase and NP molecules to the viral genome segments precludes RIG-I binding. svRNAs do not affect the innate immune responses during viral infections, likely because they are single-stranded and/or readily bound by the viral RNA polymerase (Perez et al. 2010). In contrast, mvRNAs and some DI RNAs are efficiently bound by RIG-I and able to induce innate immune responses (Baum et al. 2010; te Velthuis et al. 2018). Avian and pandemic influenza viruses were found to produce high levels of mvRNAs, which correlated with the activation of antiviral genes, suggesting that production of mvRNAs by unadapted avian influenza virus polymerases in mammalian cells could be a biomarker for the “cytokine storm” phenomenon that underlies the high virulence associated with infections with these viruses (Fig. 5C). However, reduced accumulation of defective viral RNA during infection of patients with pandemic 2009 viruses has been linked to severe outcome, suggesting that low levels of defective RNAs are beneficiary through activating a protective level of antiviral innate immune responses (Vasilijevic et al. 2017). The propensity of the RNA polymerase to generate subgenomic RNAs such as mvRNAs and DI RNAs is likely to be one of the critical factors in determining the magnitude of an innate immune response generated against a particular strain of influenza virus, which helps to explain previous observations showing that the RNA polymerase contributes to influenza virus virulence.

CONCLUDING REMARKS

Recent years witnessed an unprecedented progress in understanding the structure of the influenza virus RNA polymerase, providing insights into the function of the RNA synthesis machinery in transcription of viral genes, replication of the viral RNA genome, as well in determining host range and virulence. High-

resolution structures solved by using cryo-EM and crystallography offer insights into the atomic details of the initiation of transcription and replication by the influenza virus RNA polymerase. Understanding the interplay between the virus and the host and in particular the dependence of viral transcription and replication on host factors provide novel insights into the adaptation of influenza A viruses to mammalian hosts. The identification of defective viral RNAs, such as mvRNAs, and their link to innate immune activation, furthers our understanding of the molecular basis of the virulence of influenza viruses. However, many long-standing questions still remain unanswered, including how the ANP32 proteins promote the replicase function of the polymerase, how defective viral RNAs such as DI RNAs and mvRNAs are generated, and what the contribution of the defective RNAs is to disease severity in humans.

ACKNOWLEDGMENTS

E.F. is supported by United Kingdom Medical Research Council program Grant MR/R009945/1. A.J.W.t.V. is supported by joint Wellcome Trust and Royal Society Grant 206579/Z/17/Z, Isaac Newton Trust Grant 17.37(r), and the National Institutes of Health Grant R21AI147172.

This article has been made freely available online courtesy of TAUNS Laboratories.

REFERENCES

- Akkina RK, Chambers TM, Nayak DP. 1984. Expression of defective-interfering influenza virus-specific transcripts and polypeptides in infected cells. *J Virol* **51**: 395–403.
- Alnaji FG, Holmes JR, Rendon G, Vera JC, Fields CJ, Martin BE, Brooke CB. 2019. Sequencing framework for the sensitive detection and precise mapping of defective interfering particle-associated deletions across influenza A and B viruses. *J Virol* **93**. doi:10.1128/JVI.00354-19
- Arranz R, Coloma R, Chichon FJ, Conesa JJ, Carrascosa JL, Valpuesta JM, Ortin J, Martin-Benito J. 2012. The structure of native influenza virion ribonucleoproteins. *Science* **338**: 1634–1637. doi:10.1126/science.1228172
- Baker SF, Ledwith MP, Mehle A. 2018. Differential splicing of ANP32A in birds alters its ability to stimulate RNA synthesis by restricted influenza polymerase. *Cell Rep* **24**: 2581–2588.e4. doi:10.1016/j.celrep.2018.08.012
- Baum A, Sachidanandam R, García-Sastre A. 2010. Preference of RIG-I for short viral RNA molecules in infected cells revealed by next-generation sequencing. *Proc Natl Acad Sci* **107**: 16303–16308. doi:10.1073/pnas.1005077107
- Boergeling Y, Rozhdestvensky TS, Schmolke M, Resa-Infante P, Robeck T, Randau G, Wolff T, Gabriel G, Brosius J, Ludwig S. 2015. Evidence for a novel mechanism of influenza virus-induced type I interferon expression by a defective RNA-encoded protein. *PLoS Pathog* **11**: e1004924. doi:10.1371/journal.ppat.1004924
- Chang S, Sun D, Liang H, Wang J, Li J, Guo L, Wang X, Guan C, Boruah BM, Yuan L, et al. 2015. Cryo-EM structure of influenza virus RNA polymerase complex at 4.3 Å resolution. *Mol Cell* **57**: 925–935. doi:10.1016/j.molcel.2014.12.031
- Chenavas S, Estrozi LF, Slama-Schwok A, Delmas B, Di Priamo C, Baudin F, Li X, Crepin T, Ruigrok RW. 2013. Monomeric nucleoprotein of influenza A virus. *PLoS Pathog* **9**: e1003275. doi:10.1371/journal.ppat.1003275
- Coloma R, Valpuesta JM, Arranz R, Carrascosa JL, Ortin J, Martin-Benito J. 2009. The structure of a biologically active influenza virus ribonucleoprotein complex. *PLoS Pathog* **5**: e1000491. doi:10.1371/journal.ppat.1000491
- Dadonaite B, Gilberston B, Knight ML, Trifkovic S, Rockman S, Laederach A, Brown LE, Fodor E, Bauer DLV. 2019. The structure of the influenza A virus genome. *Nat Microbiol* **4**: 1781–1789. doi:10.1038/s41564-019-0513-7
- de Jong MD, Simmons CP, Thanh TT, Hien VM, Smith GJ, Chau TN, Hoang DM, Chau NV, Khanh TH, Dong VC, et al. 2006. Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med* **12**: 1203–1207. doi:10.1038/nm1477
- Deng T, Vreede FT, Brownlee GG. 2006. Different de novo initiation strategies are used by influenza virus RNA polymerase on its cRNA and viral RNA promoters during viral RNA replication. *J Virol* **80**: 2337–2348. doi:10.1128/JVI.80.5.2337-2348.2006
- De Vlugt C, Sikora D, Rocheleau L, Pelchat M. 2019. Priming and realignment by the influenza A virus RdRp is dependent on the length of the host primers and the extent of base pairing to viral RNA. *Virology* **536**: 91–100. doi:10.1016/j.virol.2019.08.002
- Dimmock NJ, Easton AJ. 2014. Defective interfering influenza virus RNAs: time to reevaluate their clinical potential as broad-spectrum antivirals? *J Virol* **88**: 5217–5227. doi:10.1128/JVI.03193-13
- Domingues P, Hale BG. 2017. Functional insights into ANP32A-dependent influenza A virus polymerase host restriction. *Cell Rep* **20**: 2538–2546. doi:10.1016/j.celrep.2017.08.061
- Eisfeld AJ, Neumann G, Kawaoka Y. 2015. At the centre: influenza A virus ribonucleoproteins. *Nat Rev Microbiol* **13**: 28–41. doi:10.1038/nrmicro3367
- Engelhardt OG, Smith M, Fodor E. 2005. Association of the influenza A virus RNA-dependent RNA polymerase with cellular RNA polymerase II. *J Virol* **79**: 5812–5818. doi:10.1128/JVI.79.9.5812-5818.2005
- Fan H, Walker AP, Carrique L, Keown JR, Serna Martin I, Karia D, Sharps J, Hengrung N, Pardon E, Steyaert J, et al. 2019. Structures of influenza A virus RNA polymerase offer insight into viral genome replication. *Nature* **573**: 287–290. doi:10.1038/s41586-019-1530-7

E. Fodor and A.J.W. te Velthuis

- Ferhadian D, Contrant M, Printz-Schweigert A, Smyth RP, Paillart JC, Marquet R. 2018. Structural and functional motifs in influenza virus RNAs. *Front Microbiol* **9**: 559. doi:10.3389/fmicb.2018.00559
- Forero A, Tisoncik-Go J, Watanabe T, Zhong G, Hatta M, Tchitchek N, Selinger C, Chang J, Barker K, Morrison J, et al. 2016. The 1918 influenza virus PB2 protein enhances virulence through the disruption of inflammatory and Wnt-mediated signaling in mice. *J Virol* **90**: 2240–2253. doi:10.1128/JVI.02974-15
- Gallagher JR, Torian U, McCraw DM, Harris AK. 2017. Structural studies of influenza virus RNPs by electron microscopy indicate molecular contortions within NP supra-structures. *J Struct Biol* **197**: 294–307. doi:10.1016/j.jmb.2016.12.007
- Hengrung N, El Omari K, Serna Martin I, Vreede FT, Cusack S, Rambo RP, Vonnrhein C, Bricogne G, Stuart DI, Grimes JM, et al. 2015. Crystal structure of the RNA-dependent RNA polymerase from influenza C virus. *Nature* **527**: 114–117. doi:10.1038/nature15525
- Hutchinson EC, Denham EM, Thomas B, Trudgian DC, Hester SS, Ridlova G, York A, Turrell L, Fodor E. 2012. Mapping the phosphoproteome of influenza A and B viruses by mass spectrometry. *PLoS Pathog* **8**: e1002993. doi:10.1371/journal.ppat.1002993
- Jennings PA, Finch JT, Winter G, Robertson JS. 1983. Does the higher order structure of the influenza virus ribonucleoprotein guide sequence rearrangements in influenza viral RNA? *Cell* **34**: 619–627. doi:10.1016/0092-8674(83)90394-X
- Jorba N, Area E, Ortin J. 2008. Oligomerization of the influenza virus polymerase complex in vivo. *J Gen Virol* **89**: 520–524. doi:10.1099/vir.0.83387-0
- Kash JC, Tumpsey TM, Proll SC, Carter V, Perwitasari O, Thomas MJ, Basler CF, Palese P, Taubenberger JK, García-Sastre A, et al. 2006. Genomic analysis of increased host immune and cell death responses induced by 1918 influenza virus. *Nature* **443**: 578–581. doi:10.1038/nature05181
- Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, Uematsu S, Jung A, Kawai T, Ishii KJ, et al. 2006. Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature* **441**: 101–105. doi:10.1038/nature04734
- Klinkhamer J, Schnepf D, Ye L, Schwaderlapp M, Gad HH, Hartmann R, Garcin D, Mahlaköiv T, Staeheli P. 2018. IFN- λ prevents influenza virus spread from the upper airways to the lungs and limits virus transmission. *eLife* **7**: e33354. doi:10.7554/eLife.33354
- Koire A, Gilbert BE, Sugang R, Zechiedrich L. 2016. Repurposed transcriptomic data reveal small viral RNA produced by influenza virus during infection in mice. *PLoS ONE* **11**: e0165729. doi:10.1371/journal.pone.0165729
- Kouba T, Drncova P, Cusack S. 2019. Structural snapshots of actively transcribing influenza polymerase. *Nat Struct Mol Biol* **26**: 460–470. doi:10.1038/s41594-019-0232-z
- Krammer F, Smith GJD, Fouchier RAM, Peiris M, Kedzieriska K, Doherty PC, Palese P, Shaw ML, Treanor J, Webster RG, et al. 2018. Influenza. *Nat Rev Dis Primers* **4**: 3. doi:10.1038/s41572-018-0002-y
- Lee N, Le Sage V, Nanni AV, Snyder DJ, Cooper VS, Lakdawala SS. 2017. Genome-wide analysis of influenza viral RNA and nucleoprotein association. *Nucleic Acids Res* **45**: 8968–8977. doi:10.1093/nar/gkx584
- Li H, Bradley KC, Long JS, Frise R, Ashcroft JW, Hartgroves LC, Shelton H, Makris S, Johansson C, Cao B, et al. 2018. Internal genes of a highly pathogenic H5N1 influenza virus determine high viral replication in myeloid cells and severe outcome of infection in mice. *PLoS Pathog* **14**: e1006821. doi:10.1371/journal.ppat.1006821
- Liu G, Lu Y, Liu Q, Zhou Y. 2019. Inhibition of ongoing influenza A virus replication reveals different mechanisms of RIG-I activation. *J Virol* **93**. doi:10.1128/JVI.02066-18
- Lo CY, Tang YS, Shaw PC. 2018. Structure and function of influenza virus ribonucleoprotein. *Subcell Biochem* **88**: 95–128. doi:10.1007/978-981-10-8456-0_5
- Long JS, Giotis ES, Moncorgé O, Frise R, Mistry B, James J, Morisson M, Iqbal M, Vignal A, Skinner MA, et al. 2016. Species difference in ANP32A underlies influenza A virus polymerase host restriction. *Nature* **529**: 101–104. doi:10.1038/nature16474
- Lukarska M, Fournier G, Pflug A, Resa-Infante P, Reich S, Naffakh N, Cusack S. 2017. Structural basis of an essential interaction between influenza polymerase and Pol II CTD. *Nature* **541**: 117–121. doi:10.1038/nature20594
- Martínez-Alonso M, Hengrung N, Fodor E. 2016. RNA-free and ribonucleoprotein-associated influenza virus polymerases directly bind the serine-5-phosphorylated carboxyl-terminal domain of host RNA polymerase II. *J Virol* **90**: 6014–6021. doi:10.1128/JVI.00494-16
- Mehle A, Doudna JA. 2009. Adaptive strategies of the influenza virus polymerase for replication in humans. *Proc Natl Acad Sci* **106**: 21312–21316. doi:10.1073/pnas.0911915106
- Moeller A, Kirchdoerfer RN, Potter CS, Carragher B, Wilson IA. 2012. Organization of the influenza virus replication machinery. *Science* **338**: 1631–1634. doi:10.1126/science.1227270
- Mondal A, Potts GK, Dawson AR, Coon JJ, Mehle A. 2015. Phosphorylation at the homotypic interface regulates nucleoprotein oligomerization and assembly of the influenza virus replication machinery. *PLoS Pathog* **11**: e1004826. doi:10.1371/journal.ppat.1004826
- Mondal A, Dawson AR, Potts GK, Freiburger EC, Baker SF, Moser LA, Bernard KA, Coon JJ, Mehle A. 2017. Influenza virus recruits host protein kinase C to control assembly and activity of its replication machinery. *eLife* **6**: e26910. doi:10.7554/eLife.26910
- Ng AK, Zhang H, Tan K, Li Z, Liu JH, Chan PK, Li SM, Chan WY, Au SW, Joachimiak A, et al. 2008. Structure of the influenza virus A H5N1 nucleoprotein: implications for RNA binding, oligomerization, and vaccine design. *FASEB J* **22**: 3638–3647. doi:10.1096/fj.08-112110
- Ortega J, Martín-Benito J, Zurcher T, Valpuesta JM, Carrascosa JL, Ortin J. 2000. Ultrastructural and functional analyses of recombinant influenza virus ribonucleoproteins suggest dimerization of nucleoprotein during virus amplification. *J Virol* **74**: 156–163. doi:10.1128/JVI.74.1.156-163.2000
- Oymans J, te Velthuis AJW. 2018. A mechanism for priming and realignment during influenza A virus replication. *J Virol* **92**: e01773.

- Peacock TP, Sheppard CM, Staller E, Barclay WS. 2019. Host determinants of influenza RNA synthesis. *Annu Rev Virol* **6**: 215–233. doi:10.1146/annurev-virology-092917-043339
- Peng Q, Liu Y, Peng R, Wang M, Yang W, Song H, Chen Y, Liu S, Han M, Zhang X, et al. 2019. Structural insight into RNA synthesis by influenza D polymerase. *Nat Microbiol* **4**: 1750–1759. doi:10.1038/s41564-019-0487-5
- Perez JT, Varble A, Sachidanandam R, Zlatev I, Manoharan M, García-Sastre A, tenOever BR. 2010. Influenza A virus-generated small RNAs regulate the switch from transcription to replication. *Proc Natl Acad Sci* **107**: 11525–11530. doi:10.1073/pnas.1001984107
- Perez JT, Zlatev I, Aggarwal S, Subramanian S, Sachidanandam R, Kim B, Manoharan M, Tenover BR. 2012. A small-RNA enhancer of viral polymerase activity. *J Virol* **86**: 13475–13485. doi:10.1128/JVI.02295-12
- Pflug A, Guilligay D, Reich S, Cusack S. 2014. Structure of influenza A polymerase bound to the viral RNA promoter. *Nature* **516**: 355–360. doi:10.1038/nature14008
- Pflug A, Lukarska M, Resa-Infante P, Reich S, Cusack S. 2017. Structural insights into RNA synthesis by the influenza virus transcription-replication machine. *Virus Res* **234**: 103–117. doi:10.1016/j.virusres.2017.01.013
- Pflug A, Gaudon S, Resa-Infante P, Lethier M, Reich S, Schulze WM, Cusack S. 2018. Capped RNA primer binding to influenza polymerase and implications for the mechanism of cap-binding inhibitors. *Nucleic Acids Res* **46**: 956–971. doi:10.1093/nar/gkx1210
- Pichlmair A, Schulz O, Tan CP, Naslund TI, Liljestrom P, Weber F, Reis e Sousa C. 2006. RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-phosphates. *Science* **314**: 997–1001. doi:10.1126/science.1132998
- Plotch SJ, Bouloy M, Ulmanen I, Krug RM. 1981. A unique cap(m7GpppXm)-dependent influenza virion endonuclease cleaves capped RNAs to generate the primers that initiate viral RNA transcription. *Cell* **23**: 847–858. doi:10.1016/0092-8674(81)90449-9
- Poon LL, Pritlove DC, Fodor E, Brownlee GG. 1999. Direct evidence that the poly(A) tail of influenza A virus mRNA is synthesized by reiterative copying of a U track in the virion RNA template. *J Virol* **73**: 3473–3476.
- Rehwinkel J, Tan CP, Goubau D, Schulz O, Pichlmair A, Bier K, Robb N, Vreede F, Barclay W, Fodor E, et al. 2010. RIG-I detects viral genomic RNA during negative-strand RNA virus infection. *Cell* **140**: 397–408. doi:10.1016/j.cell.2010.01.020
- Reich S, Guilligay D, Pflug A, Malet H, Berger I, Crepin T, Hart D, Lunardi T, Nanao M, Ruigrok RW, et al. 2014. Structural insight into cap-snatching and RNA synthesis by influenza polymerase. *Nature* **516**: 361–366. doi:10.1038/nature14009
- Robb NC, te Velthuis AJ, Wieneke R, Tampe R, Cordes T, Fodor E, Kapanidis AN. 2016. Single-molecule FRET reveals the pre-initiation and initiation conformations of influenza virus promoter RNA. *Nucleic Acids Res* **44**: 10304–10315.
- Robb NC, te Velthuis AJW, Fodor E, Kapanidis AN. 2019. Real-time analysis of single influenza virus replication complexes reveals large promoter-dependent differences in initiation dynamics. *Nucleic Acids Res* **47**: 6466–6477. doi:10.1093/nar/gkz313
- Robertson JS, Schubert M, Lazzarini RA. 1981. Polyadenylation sites for influenza virus mRNA. *J Virol* **38**: 157–163.
- Russell AB, Elshina E, Kowalsky JR, te Velthuis AJW, Bloom JD. 2019. Single-cell virus sequencing of influenza infections that trigger innate immunity. *J Virol* **93**. doi:10.1128/JVI.00500-19
- Saira K, Lin X, DePasse JV, Halpin R, Twaddle A, Stockwell T, Angus B, Cozzi-Lepri A, Delfino M, Dugan V, et al. 2013. Sequence analysis of in vivo defective interfering-like RNA of influenza A H1N1 pandemic virus. *J Virol* **87**: 8064–8074. doi:10.1128/JVI.00240-13
- Serna Martin I, Hengrune N, Renner M, Sharps J, Martínez-Alonso M, Masiulis S, Grimes JM, Fodor E. 2018. A mechanism for the activation of the influenza virus transcriptase. *Mol Cell* **70**: 1101–1110.e4. doi:10.1016/j.molcel.2018.05.011
- Staller E, Sheppard CM, Neasham PJ, Mistry B, Peacock TP, Goldhill DH, Long JS, Barclay WS. 2019. ANP32 proteins are essential for influenza virus replication in human cells. *J Virol* **93**: e00217. doi:10.1128/JVI.00217-19
- Subbarao EK, London W, Murphy BR. 1993. A single amino acid in the PB2 gene of influenza A virus is a determinant of host range. *J Virol* **67**: 1761–1764.
- Sugiyama K, Kawaguchi A, Okuwaki M, Nagata K. 2015. pp32 and APRIL are host cell-derived regulators of influenza virus RNA synthesis from cRNA. *eLife* **4**: e08939. doi:10.7554/eLife.08939
- Taubenberger JK, Kash JC, Morens DM. 2019. The 1918 influenza pandemic: 100 years of questions answered and unanswered. *Sci Transl Med* **11**: eaau5485. doi:10.1126/scitranslmed.aau5485
- te Velthuis AJ. 2014. Common and unique features of viral RNA-dependent polymerases. *Cell Mol Life Sci* **71**: 4403–4420. doi:10.1007/s00018-014-1695-z
- te Velthuis AJ, Fodor E. 2016. Influenza virus RNA polymerase: Insights into the mechanisms of viral RNA synthesis. *Nat Rev Microbiol* **14**: 479–493. doi:10.1038/nrmicro.2016.87
- te Velthuis AJW, Oymans J. 2018. Initiation, elongation, and realignment during influenza virus mRNA synthesis. *J Virol* **92**: e01775-17.
- te Velthuis AJ, Robb NC, Kapanidis AN, Fodor E. 2016. The role of the priming loop in influenza A virus RNA synthesis. *Nat Microbiol* **1**: 16029. doi:10.1038/nmicrobiol.2016.29
- te Velthuis AJW, Long JC, Bauer DLV, Fan RLY, Yen HL, Sharps J, Siegers JY, Killip MJ, French H, Oliva-Martin MJ, et al. 2018. Mini viral RNAs act as innate immune agonists during influenza virus infection. *Nat Microbiol* **3**: 1234–1242. doi:10.1038/s41564-018-0240-5
- Thierry E, Guilligay D, Kosinski J, Bock T, Gaudon S, Round A, Pflug A, Hengrune N, El Omari K, Baudin F, et al. 2016. Influenza polymerase can adopt an alternative configuration involving a radical repacking of PB2 domains. *Mol Cell* **61**: 125–137. doi:10.1016/j.molcel.2015.11.016
- Turrell L, Lyall JW, Tiley LS, Fodor E, Vreede FT. 2013. The role and assembly mechanism of nucleoprotein in influenza A virus ribonucleoprotein complexes. *Nat Commun* **4**: 1591. doi:10.1038/ncomms2589
- Turrell L, Hutchinson EC, Vreede FT, Fodor E. 2015. Regulation of influenza A virus nucleoprotein oligomerization

E. Fodor and A.J.W. te Velthuis

- by phosphorylation. *J Virol* **89**: 1452–1455. doi:10.1128/JVI.02332-14
- Vasilijevic J, Zamarreño N, Oliveros JC, Rodriguez-Frandsen A, Gómez G, Rodriguez G, Pérez-Ruiz M, Rey S, Barba I, Pozo F, et al. 2017. Reduced accumulation of defective viral genomes contributes to severe outcome in influenza virus infected patients. *PLoS Pathog* **13**: e1006650. doi:10.1371/journal.ppat.1006650
- Vreede FT, Jung TE, Brownlee GG. 2004. Model suggesting that replication of influenza virus is regulated by stabilization of replicative intermediates. *J Virol* **78**: 9568–9572. doi:10.1128/JVI.78.17.9568-9572.2004
- Walker AP, Fodor E. 2019. Interplay between influenza virus and the host RNA polymerase II transcriptional machinery. *Trends Microbiol* **27**: 398–407. doi:10.1016/j.tim.2018.12.013
- Williams GD, Townsend D, Wylie KM, Kim PJ, Amarasinghe GK, Kutluay SB, Boon ACM. 2018. Nucleotide resolution mapping of influenza A virus nucleoprotein-RNA interactions reveals RNA features required for replication. *Nat Commun* **9**: 465. doi:10.1038/s41467-018-02886-w
- Yamada S, Hatta M, Staker BL, Watanabe S, Imai M, Shinya K, Sakai-Tagawa Y, Ito M, Ozawa M, Watanabe T, et al. 2010. Biological and structural characterization of a host-adapting amino acid in influenza virus. *PLoS Pathog* **6**: e1001034. doi:10.1371/journal.ppat.1001034
- Ye Q, Krug RM, Tao YJ. 2006. The mechanism by which influenza A virus nucleoprotein forms oligomers and binds RNA. *Nature* **444**: 1078–1082. doi:10.1038/nature05379
- Yoneyama M, Kikuchi M, Natsukawa T, Shinobu N, Imai-zumi T, Miyagishi M, Taira K, Akira S, Fujita T. 2004. The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat Immunol* **5**: 730–737. doi:10.1038/ni1087
- York A, Hengrung N, Vreede FT, Huiskonen JT, Fodor E. 2013. Isolation and characterization of the positive-sense replicative intermediate of a negative-strand RNA virus. *Proc Natl Acad Sci* **110**: E4238–E4245. doi:10.1073/pnas.1315068110
- Zhang H, Zhang Z, Wang Y, Wang M, Wang X, Zhang X, Ji S, Du C, Chen H, Wang X. 2019. Fundamental contribution and host range determination of ANP32A and ANP32B in influenza A virus polymerase activity. *J Virol* **93**. doi:10.1128/JVI.00174-19

Structure and Function of the Influenza Virus Transcription and Replication Machinery

Ervin Fodor and Aartjan J.W. te Velthuis

Cold Spring Harb Perspect Med 2020; doi: 10.1101/cshperspect.a038398 originally published online December 23, 2019

Subject Collection [Influenza: The Cutting Edge](#)

Emerging HxNy Influenza A Viruses

William J. Liu, Yan Wu, Yuhai Bi, et al.

Equine Influenza

Thomas M. Chambers

Human Influenza Epidemiology

Sukhyun Ryu and Benjamin J. Cowling

Host Cell Factors That Interact with Influenza Virus Ribonucleoproteins

Ecco Staller and Wendy S. Barclay

Induction and Evasion of Type-I Interferon Responses during Influenza A Virus Infection

Raquel Muñoz-Moreno, Carles Martínez-Romero and Adolfo García-Sastre

Structure and Function of Influenza Polymerase

Joanna M. Wandzik, Tomas Kouba and Stephen Cusack

H7N9 Influenza Virus in China

Chengjun Li and Hualan Chen

H5 Influenza Viruses in Egypt

Rabeh El-Shesheny, Ahmed Kandeil, Ahmed Mostafa, et al.

Antivirals Targeting the Neuraminidase

Larisa Gubareva and Teena Mohan

Accessory Gene Products of Influenza A Virus

Rute M. Pinto, Samantha Lycett, Eleanor Gaunt, et al.

Influenza Immunization in the Context of Preexisting Immunity

Susanne L. Linderman, Ali H. Ellebedy, Carl Davis, et al.

Hemagglutinin Structure and Activities

Steven J. Gamblin, Sébastien G. Vachieri, Xiaoli Xiong, et al.

Live Attenuated Cold-Adapted Influenza Vaccines

Kanta Subbarao

Next-Generation Influenza Vaccines

Masaru Kanekiyo and Barney S. Graham

Selective Genome Packaging Mechanisms of Influenza A Viruses

Takeshi Noda

Systems Biological Analysis of Immune Response to Influenza Vaccination

Mario Cortese, Amy C. Sherman, Nadine G. Roupheal, et al.

For additional articles in this collection, see <http://perspectivesinmedicine.cshlp.org/cgi/collection/>