The Role of Somatic Mutations in Acute Myeloid Leukemia Pathogenesis

Ashwin Kishtagari^{1,2} and Ross L. Levine^{3,4,5,6,7}

¹Department of Translational Hematology and Oncology Research, ²Department of Hematology and Oncology, Taussig Cancer Institute, Cleveland Clinic, Cleveland, Ohio 44195, USA

³Human Oncology and Pathogenesis Program, ⁴Department of Medicine, Leukemia Service,

⁵Center for Epigenetics Research, ⁶Center for Hematologic Malignancies, ⁷Molecular Cancer Medicine Service, Memorial Sloan Kettering Cancer Center, New York, New York 10065, USA

Correspondence: leviner@mskcc.org

Acute myeloid leukemia (AML) is characterized by attenuation of lineage differentiation trajectories that results in impaired hematopoiesis and enhanced self-renewal. To date, sequencing studies have provided a rich landscape of information on the somatic mutations that contribute to AML pathogenesis. These studies show that most AML genomes harbor relatively fewer mutations, which are acquired in a stepwise manner. Our understanding of the genetic basis of leukemogenesis informs a broader understanding of what initiates and maintains the AML clone and informs the development of prognostic models and mechanism-based therapeutic strategies. Here, we explore the current knowledge of genetic and epigenetic aberrations in AML pathogenesis and how recent studies are expanding our knowledge of leukemogenesis and using this to accelerate therapeutic development for AML patients.

Acute myeloid leukemia (AML) is a clonal disorder initiated in hematopoietic stem and progenitor cells (HSPCs) that is characterized by impaired myeloid differentiation and abnormal proliferation at the expense of the normal hematopoietic system. AML is the most common form of acute leukemia in adults; an estimated 19,940 people will be diagnosed in 2020, and 11,180 patients will die of the disease (Siegel et al. 2020). AML is the 10th leading

cause of cancer deaths in the United States. AML is most common in older adults, with a median age at diagnosis of 68 yr. Treatment of AML remains a significant challenge owing to the diverse molecular mechanisms contributing to different AML subtypes, and limited therapeutic options. Even though two-thirds of fit patients with newly diagnosed AML achieve morphological remission with intensive chemotherapy consisting of cytarabine and an anthra-

Editors: Michael G. Kharas, Ross L. Levine, and Ari M. Melnick

Additional Perspectives on Leukemia and Lymphoma: Molecular and Therapeutic Insights available at www.perspectivesinmedicine.org

Copyright © 2021 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a034975

Cite this article as *Cold Spring Harb Perspect Med* 2021;11:a034975

A. Kishtagari and R.L. Levine

cycline, the majority of these patients eventually relapse (Döhner et al. 2017; Kishtagari et al. 2020).

The first whole-genome sequencing of a cancer patient was of a patient with cytogenetically normal AML, reported more than a decade ago (Ley et al. 2008). Since then, gene discovery studies have identified an extensive catalog of recurrent somatic mutations in different AML subtypes (Mardis et al. 2009; Patel et al. 2012; Cancer Genome Atlas Research et al. 2013; Papaemmanuil et al. 2016; Tyner et al. 2018). These discovery studies have led to a greater understanding of AML biology, including stepwise accumulation of genetic alterations and recurrent mutations in epigenetic modifiers in AML. The types of mutations found in AML include chromosomal rearrangements (such as translocations or inversion), gain or loss of chromosomes (aneuploidy), total or partial gene deletion, point mutation, insertion, or gene duplication/amplification. AML can develop after an antecedent myeloid malignancy (secondary AML [s-AML]), after leukemogenic therapy (therapy-related AML [t-AML]), or without an identifiable prodrome or known exposure (de novo AML). The therapeutic strategy in “fit” patients with AML (primarily influenced by patient-related factors such as advanced age, performance status, and pretreatment comorbidities) is divided into induction and postremission therapy. Induction refers to the use of cytotoxic therapy (combination of cytarabine and an anthracycline) to induce remission, preferably without measurable residual disease (MRD). Standard postremission strategies include lower-intensity cytotoxic therapy and hematopoietic stem cell transplantation (HSCT). The use of different therapies in the postremission treatment is largely determined by prognostic risk stratification (Table 1). The treatment of AML patients who are “unfit” for intensive therapy include low-dose cytarabine, hypomethylating agents, and combination treatment strategies with novel agents (e.g., venetoclax, glasdegib). Recent data suggests remarkable efficacy for venetoclax/azacytidine in older adults with AML who are not candidates for intensive therapy, such that this regimen is quickly emerg-

ing as a new alternative for first-line therapy for older adults with AML (Pollyea et al. 2018).

In this review, we provide an overview of AML pathophysiology through the prism of genomic and epigenomic alterations and discuss how these mechanisms are perturbed in leukemogenesis. Acute promyelocytic leukemia (APL), a distinct subtype of AML, will not be addressed in this review, as we would posit it is a unique malignancy with distinct mechanisms of transformation and therapeutic options. In essence, we will discuss the pathogenesis of AML by addressing the role of recurrent cytogenetic abnormalities in AML pathogenesis, the underlying molecular events that contribute to AML, how our evolving understanding of the molecular basis of AML is refining prognostic schema, and how these insights are leading to new therapeutic approaches.

CHROMOSOMAL ABNORMALITIES

In AML, detection of chromosomal abnormalities by cytogenetic analysis is critically important for diagnosis, risk stratification, and therapeutic decision-making (Fig. 1). Chromosomal abnormalities are detected in ~50% of patients with AML at the time of diagnosis. Chromosomal abnormalities most commonly contribute to leukemogenesis through the production of a fusion oncoprotein and/or by altering *cis*-regulatory elements. These large-scale copy-number alterations can serve as founding events and drive disease evolution in AML. Detection of t(8;21)(q22;q22.1), inv(16)(p13.1;q22), t(16;16)(p13.1;q22), or translocations generating *PML-RARA* fusion transcripts (APL) allow the diagnosis of AML, regardless of the percentage of myeloblasts in the bone marrow or peripheral blood.

Core Binding Factor Rearrangements

The inv(16)(p13;q22) or t(16;16)(p13.1;q22) and t(8;21)(q22;q22.1) translocations lead to the expression of the *CBFB-MYH11* and *RUNX1-RUNX1T1* fusion oncogenes, respectively. In steady state hematopoiesis, core bind-

Table 1. The European LeukemiaNET (ELN) 2017 risk stratification of acute myeloid leukemia (AML)

Risk category	Genetic abnormality
Favorable	<p>t(8;21)(q22;q22.1); <i>RUNX1-RUNX1T1</i></p> <p>inv(16)(p13.1;q22) or t(16;16)(p13.1;q22); <i>CBFB-MYH11</i></p> <p>Mutated <i>NPM1</i> without <i>FLT3-ITD</i> or with <i>FLT3-ITD</i>^{low} = allelic ratio < 0.5</p> <p>Biallelic mutated <i>CEBPA</i></p>
Intermediate	<p>Mutated <i>NPM1</i> and <i>FLT3-ITD</i>^{high} = allelic ratio > 0.5</p> <p>Wild-type <i>NPM1</i> without <i>FLT3-ITD</i> or with <i>FLT3-ITD</i>^{low} (without adverse-risk genetic lesions)</p> <p>t(9;11)(p21.3;q23.3); <i>MLLT3-KMT2A</i></p> <p>Cytogenetic abnormalities not classified as favorable or adverse</p>
Adverse	<p>t(6;9)(p23;q34.1); <i>DEK-NUP214</i></p> <p>t(v;11q23.3); <i>KMT2A</i> rearranged</p> <p>t(9;22)(q34.1;q11.2); <i>BCR-ABL1</i></p> <p>inv(3)(q21.3;q26.2) or t(3;3)(q21.3;q26.2); <i>GATA2,MECOM(EVI1)</i></p> <p>–5 or del(5q); –7; –17/abn(17p)</p> <p>Complex karyotype, monosomal karyotype</p> <p>Wild-type <i>NPM1</i> and <i>FLT3-ITD</i>^{high}</p> <p>Mutated <i>RUNX1</i></p> <p>Mutated <i>ASXL1</i></p> <p>Mutated <i>TP53</i></p>

Data reprinted from Döhner et al. (2017), with permission from The American Society of Hematology, © 2017.

ing factor beta (CBFβ) and RUNX1 heterodimerize to bind DNA and recruit lineage-defining transcription factors to regulate hematopoietic differentiation (Tahirov et al. 2001). In CBF-AML, chromosomal rearrangements alter this transcriptional complex, thus dysregulating normal hematopoiesis. These rearrangements are present in ~15% of AML cases. They are the most responsive AML subtypes to intensive chemotherapy, including postremission therapy with high-dose cytarabine (HiDAC) such that patients with these fusion oncoproteins are treated with induction chemotherapy followed by HiDAC consolidation as standard of care (Bloomfield et al. 1998; Byrd et al. 2004). Patients with CBF-AML usually

have a relatively good outcome with overall survival rates of 60%–70% in adults. However, the expression of CD56 and the activating mutations in *KIT* have been correlated to greater risk of recurrence and shorter survival. The *KIT* mutations are found in 20%–25% of t(8;21)(q22;q22) and in ~30% of inv(16) (p13;q22) patients (Paschka et al. 2006). However, the role of *KIT* mutations in prognostication and in molecularly targeted therapies for AML requires further prognostic and therapeutic studies. Both *CBFB-MYH11* and *RUNX1-RUNX1T1* fusion transcripts are well-established markers for MRD monitoring by real-time quantitative polymerase chain reaction (qRT-PCR) to guide postremission therapy (Schuurhuis et al. 2018).

A. Kishtagari and R.L. Levine

Figure 1. Overview of frequently mutated cellular pathways in acute myeloid leukemia (AML). These include (1) *CBFB-MYH11* and *RUNX1-RUNX1T1* fusion oncogenes; (2) nucleophosmin 1 (*NPM1*); (3) transcription factors *RUNX1*, *CEBPA*, and *GATA2*; (4) tumor suppressors *TP53*; (5) signal transduction, including *FLT3-ITD*, *FLT3-TKD*, *KIT*, *CBL*, *NRAS*, *KRAS*, *PTPN11*, and *NF1*; and (6) RNA splicing: *SF3B1*, *SRSF2*, *U2AF1*, and *ZRSR2*. Mutated genes are highlighted with a red star. (Refer to the main text for detailed explanations of the depicted processes.) (Figure reprinted with permission, Cleveland Clinic Center for Medical Art & Photography © 2019–2020. All rights reserved.)

KMT2A (MLL1) Rearrangements

The mixed-lineage leukemia gene (*KMT2A*, also known as *MLL1*) encodes a histone H3 lysine 4 (H3K4) methyltransferase that regulates the expression of target genes, including homeobox (*Hox*) genes. In *MLL1* fusions, *MLL1* retains the DNA-binding amino-terminal domain, and the catalytic methyltransferase activity carboxy-terminal SET domain is replaced by a fusion partner of one of more than 70 partner genes, most commonly *AF4* t(4;11), *AF9* t(9;11), *ENL* t(11;19), *AF10* t(10;11), and *ELL* t(11;19)

(Krivtsov and Armstrong 2007). The most common fusion partners of *MLL1* are members of the DOT1L complex and/or the super elongation complex (SEC), which are both involved in transcriptional control. *MLL1*-fusion leukemias are characterized by increased expression of *HOXA9*, *MEIS1*, and *MEF2C*, which encode for leukemogenic transcription factors (Armstrong et al. 2002). *MLL1* fusions occur in as many as 10% of patients with AML and are common in t-AML following exposure to topoisomerase II inhibitors (Strissel et al. 1998). *MLL1*-fusion AML carries a poor prognosis be-

cause of refractoriness to chemotherapy and shorter period to relapse (Meyer et al. 2018). Aberrant recruitment of the DOT1L by *MLL1*-fusion leukemias provided a rationale for development of small-molecule inhibitors (e.g., pinometostat) targeting DOT1L with modest results (Stein et al. 2018). Moreover, recent data showing that *MLL1* fusions, but not wild-type *MLL1*, requiring menin as a co-factor to activate downstream target gene expression has led to the development of small-molecule inhibitors of the *MLL1*/menin interaction, which are now entering clinical trials (Yokoyama et al. 2005; Grembecka et al. 2012; Borkin et al. 2015; Krivtsov et al. 2019; Klossowski et al. 2020). Another mechanism by which *MLL1* plays a role in leukemogenesis is through an in-frame partial tandem duplication (*MLL1-PTD*) spanning exons 3 to 9, exons 3 to 10, or exons 3 to 11 (Steudel et al. 2003). *MLL1-PTDs* occur in 3.2%–11% of adult AML and more frequently present in AML with normal karyotype and trisomy 11 (Basecke et al. 2006). They confer a worse prognosis with shortened overall survival. *MLL1-PTD* AML has a distinct gene expression signature, and concomitant *DNMT3A* and *NRAS* mutations were associated with adverse clinical outcomes (Hinai et al. 2019).

Rare Translocations and Chromosomal Alterations

inv(3)(q21.3;q26.2) or *t(3;3)(q21.3;q26.2); GATA2, MECOM(EV11)*

These rearrangements result in overexpression of the proto-oncogene *MECOM* (*EV11*) at 3q26.2, through juxtaposition of the *EV11* gene with a distal *GATA2* enhancer, and simultaneously confers *GATA2* haploinsufficiency (Gröschel et al. 2014). It is a distinct entity, which is present in de novo AML or AML arising from prior myelodysplastic syndromes (MDS), and is associated with aggressive disease with minimal response to standard therapies and short survival (Lugthart et al. 2010; Rogers et al. 2014). These rearrangements occur in 1%–2% of all AML cases. Mutations in genes activating RAS/receptor tyrosine kinase signaling pathways

Somatic Mutations in Acute Myeloid Leukemia

are reported in a significant proportion of the cases with these rearrangements; this may provide a target for a rational treatment strategy in this adverse-risk patient group (Gröschel et al. 2015).

t(6;9)(p23;q34.1); DEK-NUP214

This rearrangement results in a fusion of oncogene *DEK* on chromosome 6 with the nucleoporin gene *NUP214* on chromosome 9. The resulting nucleoporin chimeric fusion protein interacts with exportin-1 (XPO1)/CRM1 and may affect the export of nuclear proteins (Saito et al. 2016). It is a distinct entity reported in ~2% of AML cases and is associated with a very poor prognosis. The majority of the patients with these rearrangements have a concomitant mutation in *FLT3-ITD*. Outcomes are dismal with standard chemotherapy with the only chance of long-term remission achieved for patients who can undergo allogeneic hematopoietic stem cell transplantation in first complete remission (Díaz-Beyá et al. 2020; Kayser et al. 2020). Given the fusion chimeric protein interaction with XPO1/CRM1, it will be interesting to see whether this fusion imparts sensitivity to XPO1 inhibitors (e.g., selinexor).

Complex Karyotype, -5/5q, -7/7q, -17/17p, +8/8q

Somatically acquired chromosome copy or segment gains, chromosomal monosomies, as well as the accumulation of karyotypic abnormalities, classified as complex karyotype, defined as three or more chromosomal abnormalities, are reported in ~5%–10% of patients with AML. These alterations carry an adverse prognosis and are associated with a poor response to cytotoxic chemotherapy. Common abnormalities include those involving chromosomes 5 and 7, particularly in AML arising out of the background of MDS, and in t-AML arising in patients treated with alkylating agents and/or radiation therapy for a previous cancer. About 50% of patients with complex karyotype abnormalities have mutations in *TP53*, and it is hypothesized that *TP53* inactivation allows cells to

A. Kishtagari and R.L. Levine

tolerate and persist in a setting of ongoing chromosomal instability. Moreover, it remains unclear whether the poor prognosis of complex karyotype AML is due to chromosomal alterations, inactivation of the p53 pathway, or both molecular events acting in concert. Haploinsufficiency of the following genes have been implicated in myeloid malignancies with monosomy 5 or large deletions of 5q [-5/del(5q)], including *RPS14*, *APC*, *CTNNA1*, *HSPA9*, *EGR1*, and *CSNK1A1* (Joslin et al. 2007; Liu et al. 2007; Ebert et al. 2008; Wang et al. 2010; Chen et al. 2011). Haploinsufficiency or loss of heterozygosity of candidate genes involved in -7/del(7q) include *SAMD9/9L*, *DOCK4*, *EZH2*, *CUX1*, *MLL3*, and *LUC7L2*, each of which has been suggested to contribute to myeloid transformation (Zhou et al. 2011; McNerney et al. 2013; Nagamachi et al. 2013; Chen et al. 2014; Hosono et al. 2014). Loss of chromosome 17p, including the *TP53* locus at 17p13.1, is associated with complex karyotype as well as abnormalities in chromosomes 5 and 7. In contrast to AML with balanced translocations, leukemias that develop in the context of 17p alterations are characterized by greater genomic instability (Liu et al. 2016). It is important to note that *NF1* is also on chromosome 17 and that some of the events involving 17q result in haploinsufficiency/loss of *NF1* and augmented MAPK signaling (Parkin et al. 2010). Other rare rearrangements, such as t(3;17)(q26;q22), involving fusion of the *EVII* gene located on chromosome 3q26.2 and *MSI2* (MUSASHI [MSI] family of RNA binding protein) located on 17q are found in AML (De Weer et al. 2008). Somatic acquisition of trisomy 8 is seen in ~10% of patients with AML. They contribute to leukemogenesis, at least in part, via amplification of *MYC*, which is located at chromosome 8p24 and is implicated in a spectrum of malignancies including AML (Sloand et al. 2007).

RECURRENT MUTATIONS IN AML

NPM1

More than 50% of patients with normal karyotype AML harbor mutations in nucleophosmin

1 (*NPM1*) (Fig. 1). *NPM1* is a multifunctional nucleus and nucleolar phosphoprotein involved in pleiotropic cellular functions (Falini et al. 2005). *NPM1* gene mutations typically occur as insertions in exon 12, causing a frameshift mutation with an added nuclear export signal (NES) motif at the carboxy-terminus; *NPM1c* protein product thus accumulates aberrantly in the cytoplasm (Falini et al. 2005; Sportoletti et al. 2015). Multiple recent publications have demonstrated that mutant *NPM1* is aberrantly localized in the cytoplasm (*NPM1c*) and this mislocalization is essential for maintaining active chromatin marks that may help maintain expression of key target genes including *HOXA* and *HOXB* genes and *MEIS1*, thereby inhibiting differentiation of leukemic cells (Brunetti et al. 2018; Gu et al. 2018). It has been suggested that *NPM1c* shuttles PU.1 (SPI1) into the cytoplasm, which abrogates PU.1 mediated repression of *HOX/MEIS1* transcription (Gu et al. 2018). The World Health Organization (WHO) classification of hematopoietic tumors recognizes *NPM1*-mutant AML as a distinct entity (Table 1) (Arber et al. 2016; Döhner et al. 2017). Strikingly, the majority of *NPM1*-mutated AML have mutations in *DNMT3A* and *FLT3*. AML patients with *NPM1* mutations without mutations in *DNMT3A* and *FLT3* carry a favorable prognosis, but the co-mutations *FLT3-ITD* and *DNMT3A* confer an adverse prognosis (Döhner et al. 2017). In *NPM1*-mutant AML, prediction of relapse risk by MRD assessment by qRT-PCR demonstrated that persistent *NPM1* positivity in blood was the sole predictor of relapse in multivariate analysis irrespective of the co-mutational status (Ivey et al. 2016; Dillon et al. 2020). This further supports its inclusion into daily clinical practice to better guide postremission therapy (Schuurhuis et al. 2018). *NPM1* mutation is considered to be a leukemia-initiating event and thus represents an ideal therapeutic target. In a recent study, using *Npm1c/Dnmt3a* mutant knock-in mice and cell lines, a model of AML development, showed *NPM1*-mutant induced self-renewal properties in myeloid progenitor cells, which act as leukemia-initiating cells. In mice, oral administration of the small molecule inhibitor,

which targets the interaction between the histone methyltransferase MLL1 and adaptor protein menin, eradicated preleukemic *Npm1c*-mutant myeloid progenitor cells, prevented AML development, and extended survival. This study demonstrates that the elimination of at-risk preleukemic cells using targeted inhibitors may be a promising strategy (Uckelmann et al. 2020).

Mutations Disrupting Transcription Factor Function in AML (Fig. 1)

Transcription factors tightly regulate normal hematopoiesis. Abrogation of specific transcription factor activity results in block in differentiation of hematopoietic progenitor cells and the development of hematologic malignancies, including AML.

RUNX1

RUNX1 encodes the sequence-specific master hematopoietic transcription factor Runt-related transcription factor 1 (*RUNX1*), which is an essential regulator of hematopoiesis by maintaining the balance between cell lineage specification, self-renewal, and proliferation (Ito et al. 2015). *RUNX1* has a highly conserved “Runt” homology domain (RHD), the motif responsible for heterodimerization with CBF β , which facilitates the specificity and affinity of *RUNX1* binding to target genes (Crute et al. 1996). More than 50 mutations in *RUNX1*, both point mutations and chromosomal rearrangements, have been reported in various hematologic malignancies, including AML. Loss-of-function somatic mutations in *RUNX1*, which are mechanistically distinct from rearrangements, occur in ~5%–15% of all patients with AML and confer a poor prognosis (Gaidzik et al. 2016). Mutations in *RUNX1* are enriched in patients with intermediate-risk (including normal karyotype AML) disease (Patel et al. 2012; Papaemmanuil et al. 2016). *RUNX1* mutations are usually secondary events that drive disease progression, and they are mutually exclusive with *NPM1* and *CEBPA* mutations (Hirsch et al. 2016; Papaemmanuil et al. 2016). They are associated

Somatic Mutations in Acute Myeloid Leukemia

with older age and lower complete response (CR) rates and inferior overall survival (OS) (Mendler et al. 2012). Germline mutations in *RUNX1* are associated with familial platelet disorder with propensity to myeloid malignancy (FPDMM), characterized by thrombocytopenia, platelet functional/ultrastructural defects, and a 20%–60% rate of transformation to overt myeloid malignancies (Song et al. 1999).

CEBPA

The transcription factor CCAAT/enhancer-binding protein α (*CEBPA*) is a master regulator involved in cell fate decisions, including myeloid-lineage commitment (Avellino and Delwel 2017). *CEBPA* is expressed as two isoforms: the full-length (p42) and the amino-terminally truncated isoform (p30) that lacks the full *trans*-activation potential of p42. *CEBPA* mutations in AML harbor *CEBPA* amino-terminal nonsense or frameshift mutations, resulting in the increased expression of the dominant-negative p30 isoform, diminishing p42 availability to promote myeloid differentiation. Carboxy-terminal in-frame mutations interrupt DNA binding and homodimerization properties of the proteins and often co-occur on a separate allele from amino-terminal mutations. This pattern of mutations produces p30/p30 homodimers as the functional *CEBPA* transcription factor entity in *CEBPA*-mutant AML (Pabst et al. 2001; Kirstetter et al. 2008). These loss-of-function *CEBPA* mutations are reported in 5%–15% of AML patients and occur mostly in younger patients with normal karyotype AML (Fröhling et al. 2004; Cancer Genome Atlas Research et al. 2013). AML patients with biallelic *CEBPA* mutations, with one allele harboring an amino-terminus mutation and the other allele having a mutation in the carboxyl terminus, exhibit a distinct gene expression signature and a favorable prognosis with the majority of younger patients achieving a cure with standard chemotherapy alone (Table 1) (Wouters et al. 2009; Fasan et al. 2014). Biallelic *CEBPA* mutations have a JAK-STAT pathway activation signature, and a high percentage of these patients also harbor gain-of-function mutations in *CSF3R* and increased sensitivity

A. Kishtagari and R.L. Levine

to JAK kinase inhibitors (such as ruxolitinib) *ex vivo* (Lavallée et al. 2016; Maxson et al. 2016). Germline mutations in *CEBPA* are associated with autosomal dominant familial AML with 100% penetrance (Smith et al. 2004).

GATA2

GATA2, a member of the zinc finger transcription factor family, is a critical transcriptional regulator of hematopoietic stem and progenitor cell differentiation and self-renewal (Crispino and Horwitz 2017). Homozygous *Gata2* knockout in mice are embryonically lethal because of the failure of definitive hematopoiesis (Tsai et al. 1994). Somatic mutations in *GATA2* are infrequent in AML, occurring in <5% of AML overall, and are clustered in normal karyotype AML with concurrent biallelic *CEBPA* mutations (Greif et al. 2012). These mutations include frameshifts, amino acid substitutions, insertions, and deletions scattered throughout the gene but most often clustered in the region encoding the zinc finger domains, leading to impaired DNA binding and affecting transcriptional activity. In its capacity as a tumor suppressor, heterozygous germline mutations in *GATA2* in coding or regulatory element regions cause a spectrum of hematopoietic disorders and a predisposition to MDS/AML, often with subsequent acquisition of somatic *ASXL1* mutations (Hahn et al. 2011; Ostergaard et al. 2011; West et al. 2014).

Mutations Altering Signal Transduction (Fig. 1)

Deregulation of signaling pathway components is frequently reported in AML. This is associated with pro-proliferative states by constitutive activation.

FLT3-ITD and FLT3-TKD

Mutations in FMS-like tyrosine kinase 3 (FLT3) occur in approximately one-third of all patients with newly diagnosed AML. These mutations can occur either as an in-frame internal tandem duplication within the juxtamembrane domain of the receptor (*FLT3-ITD*), seen in ~20%–25%

of AML, or as point mutations most commonly in the activation loop of the catalytic domain (*FLT3-TKD*). Both mutations lead to autoactivation of FLT3-kinase activity and activation of downstream signaling pathways, which lead to proliferation and block in differentiation (Meshinchi and Appelbaum 2009). AML patients with mutations in *FLT3-ITD* have an adverse prognosis compared to patients with *FLT3-TKD* or wild-type *FLT3*, and the majority of these patients relapse after chemotherapy and allogeneic HSCT (Thiede et al. 2002; Sengsayadeth et al. 2012). The poor prognosis is particularly influenced by co-mutation status (*DNMT3A* mutations, absence of *NPM1* mutations) and the allelic ratio of mutant *FLT3-ITD* to wild-type (Table 1) (Patel et al. 2012; Papaemmanuil et al. 2016). The advent of *FLT3* inhibitors (midostaurin, gilteritinib, quizartinib, and crenolanib) in various stages of clinical development represents a paradigm of targeted therapy in AML and has changed practice. These agents act through the competitive inhibition of the ATP-binding site in the FLT3 receptor; however, they vary substantially in their inhibitory properties, pharmacokinetics, and toxicity profiles. There are currently two FDA-approved small-molecule inhibitors (midostaurin and gilteritinib) of FLT3-kinase activity. In randomized clinical trial, comparing midostaurin in combination with chemotherapy or placebo with chemotherapy has been shown to significantly improve survival of untreated *FLT3*-mutant AML patients (age 18–60 yr; 74.7 vs. 25.6 mo) (Stone et al. 2017). Gilteritinib, a next-generation tyrosine kinase inhibitor, is approved for the treatment of adults who have relapsed and/or have refractory AML with *FLT3-ITD* and *FLT3-TKD* mutations (Perl et al. 2019). Of note, therapy has substantive-agent activity, whereas midostaurin has not shown significant activity when given as monotherapy.

KIT

The *KIT* gene encodes a 145-kDa transmembrane glycoprotein that is a member of the type III receptor tyrosine kinase (RTK) family (Yarden et al. 1987). Upon binding of stem cell

factor (KIT ligand), the monomeric KIT receptor dimerizes and becomes autophosphorylated at key tyrosine sites and activates diverse signal transduction cascades essential for cell proliferation/survival, self-renewal, and differentiation (Malaise et al. 2009). Somatic gain-of-function *KIT* mutations occur in 10% of AML patients, but they are clustered in patients with CBF-AML rearrangements (t(8;21)/*RUNX1-RUNX1T1*, inv(16)/*CBFB-MYH11*) and may confer a more adverse prognosis in this otherwise favorable AML subtype (Paschka et al. 2006).

CBL

Casitas B cell lymphoma (*CBL*) protein functions as E3 ligases that ubiquitinate and negatively regulate receptor tyrosine kinases, including *FLT3* and *KIT* (Mohapatra et al. 2013). Mutations in the E3 ubiquitin ligase *CBL* have been reported in ~1%–3% of AML patients. Mutations in *CBL* result in the loss of ubiquitin E3 ligase activity and resultant increased tyrosine kinase signaling by decreased proteasome-mediated degradation. In preclinical models, there is evidence that *FLT3-ITD* mutations cooperate with *CBL* to promote the rapid induction of AML (Taylor et al. 2015). Although, mutations in *CBL* cause biochemical loss of function, their functional consequences resemble canonical gain-of-function mutations (Sanada et al. 2009). The prognostic significance of these mutations in AML is unclear given the lack of robust data.

NRAS and KRAS

The Ras family of small GTPases, *NRAS* and *KRAS*, function as binary molecular switches by cycling between an active GTP-bound (RAS-GTP) and an inactive GDP-bound (RAS-GDP) confirmation that regulate downstream signaling effectors, such as Raf and PI3K (Schubert et al. 2007). Somatic gain-of-function mutations in *RAS* are among the most common oncogenic drivers in cancers leading to constitutive activation of receptor tyrosine kinases (Li et al. 2018). *NRAS* (~12%) and *KRAS* (~5%) mutations are seen in a subset of AML patients at the time of diagnosis (Bacher et al. 2006). They

are frequently acquired at the time of progression from MDS to AML and are associated with poor survival. In a recent study, the presence of *NRAS* and *KRAS* mutations in patients with AML receiving standard cytotoxic therapy was associated with decreased OS and event-free survival (EFS) (Ball et al. 2019). Mutations in epigenetic modifiers (*TET2/IDH/WT1*) often co-occur and cooperate with *NRAS* mutations to promote leukemogenesis (Papaemmanuil et al. 2016). Preclinical studies suggest this cooperativity and dependence on activated Ras signaling may be therapeutically exploited as they are preferentially sensitive to MAPK kinase (MEK) inhibition in mouse models and patient samples (Kunimoto et al. 2018). The emergence or co-occurrence of *NRAS* and *KRAS* mutant clones is a common and clinically relevant mechanism of resistance to *FLT3* inhibitors, *IDH* inhibitors, and BCL-2 inhibitors (Amatangelo et al. 2017; McMahon et al. 2019; DiNardo et al. 2020).

PTPN11

The non-receptor protein tyrosine phosphatase *PTPN11* (also known as SRC homology 2 domain-containing phosphatase 2 or SHP2) is implicated in several signaling pathways including RAS, JAK-STAT, PI3K, and others (Tartaglia et al. 2003). Mutations in *PTPN11* are present in ~5%–10% of AML cases and enriched in patients older than 60 yr with secondary AML (Makishima et al. 2017). These mutations co-occur and cooperate with *AML1-ETO* (Hatlen et al. 2016) and *MLL-AF9* (Chen et al. 2015) rearrangements.

NF1

The neurofibromin 1 gene (*NF1*), encodes neurofibromin, a GTPase-activating protein and negative regulator of Ras. Inactivating mutations in *NF1* leading to loss of neurofibromin and resulting in prolonged activation of the RAS/RAF/MAPK signaling pathway and ultimately an increased cellular proliferation (Cichowski and Jacks 2001). Recurrent somatic inactivating mutations in *NF1* have been recently reported in ~5% patients with AML (Eisfeld et al. 2018). In

A. Kishtagari and R.L. Levine

the same study, *NFI*-mutant AML patients had poor outcomes with lower complete remission rates and shorter overall survival when treated with standard chemotherapy.

Mutations in Epigenetic Modifiers: *DNMT3A*, *TET2*, *IDH1/2*, and Polycomb-Group Genes (Fig. 2)

One of the hallmarks of AML pathogenesis is disruption of epigenetic regulation. The mutations in genes encoding epigenetic modifiers are commonly acquired early and are present in the initiating clone. This has led to an increasing interest in the development of epigenetic therapies for AML.

DNMT3A

DNA methyltransferase 3A (*DNMT3A*) is a member of the DNA methyltransferase family, which includes *DNMT1*, *DNMT3A*, *DNMT3B*, and *DNMT3L* (Okano et al. 1998). It catalyzes de novo DNA methylation at the C-5 position of cytosine bases of unmethylated CpG dinucleotides. DNA methylation is a critical epigenetic modification and plays an essential role in maintaining hematopoietic homeostasis. Consistent with its function, *Dnmt3a*-null HSCs up-regulate genes implicated in HSC self-renewal and multipotency, while down-regulating expression of genes associated with HSC differentiation (Challen et al. 2011). Mutations in *DNMT3A* are observed in ~30% of AML cases.

Figure 2. Overview of frequently mutated epigenetic modifiers. This includes (1) DNA cytosine modifications *DNMT3A*, *TET2*, and *IDH1/2*; (2) histone post-translational modifications *ASXL1*, *BCOR/L1*, and *MLL1* (*KMT2A*); and (3) Cohesin complex *SMC1A*, *SMC3*, and *RAD21*, bound to *STAG1/2*. Mutated genes are highlighted with a red star. (Refer to the main text for detailed explanations of the depicted processes.) (Figure reprinted with permission, Cleveland Clinic Center for Medical Art & Photography © 2019–2020. All rights reserved.)

DNMT3A-mutant AML patients are older and most commonly present with a normal karyotype (Ley et al. 2010; Cancer Genome Atlas Research et al. 2013). The mutations in *DNMT3A* include nonsense, frameshift, and missense alterations that are enriched within the methyltransferase domain. The recurrent mutation at codon R882 is the most frequent *DNMT3A* mutation in AML patients, the resultant protein exerts a dominant-negative effect on wild-type *DNMT3A* methyltransferase activity (Holz-Schietinger et al. 2012; Cancer Genome Atlas Research et al. 2013; Russler-Germain et al. 2014). Mutations in *DNMT3A* frequently co-occur with mutations in *NPM1* and *FLT3-ITD*, which collectively confer adverse risk (Marcucci et al. 2012; Bezerra et al. 2020). *DNMT3A* mutations at codon R882 have been shown to promote anthracycline resistance through a proximal defect in nucleosome remodeling and resultant impaired DNA damaging sensing (Guryanova et al. 2016). Clonal hierarchy studies show that mutations in *DNMT3A* are one of the founder mutations shaping the course of leukemia evolution and progression (Welch et al. 2012), making this an attractive target for the development of novel therapeutic approaches. This is further underscored by studies showing that the *DNMT3A* is the most frequently mutated gene in clonal hematopoiesis (CH) (Genovese et al. 2014; Jaiswal et al. 2014; Shlush et al. 2014; Xie et al. 2014). Although essential insights into the role of *DNMT3A* mutations in leukemogenesis have emerged from human and preclinical studies, the fundamental mechanism(s) by which mutant *DNMT3A* leads to increased self-renewal in HSCs has not been delineated.

TET2

The mammalian ten-eleven-translocation (TET) family proteins TET1, TET2, and TET3, are α -ketoglutarate-dependent dioxygenases. The TET enzymes, in the presence of oxygen, reduced Fe^{2+} , and ascorbic acid, catalyze the stepwise oxidation of 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC), and 5-carboxylcytosine (5caC), with resultant loss of DNA methylation (Tahi-

liani et al. 2009; Ko et al. 2010; He et al. 2011; Ito et al. 2011). *TET2*, but not *TET1* or *TET3*, loss-of-function mutations are common across a spectrum of hematologic malignancies (Delhommeau et al. 2009; Jankowska et al. 2009). Somatic mutations in *TET2* are seen in ~20% of patients with AML, especially in patients with a prior history of MDS or myeloproliferative neoplasm (MPN) (Abdel-Wahab et al. 2009). In AML, *TET2* mutations co-occur with *NPM1*, *FLT3-ITD*, *NRAS*, and *DNMT3A* and are largely mutually exclusive with *IDH1/2* neomorphic mutations (Papaemmanuil et al. 2016; Desai et al. 2018). *TET2* mutations are among the most common mutations seen in clonal hematopoiesis (CH) (Busque et al. 2012).

Tet2^{-/-} mice develop myeloid and lymphoid neoplasms, after long latencies, through the accumulation of numerous mutations, including *Apc*, *Nf1*, *Flt3*, *Cbl*, *Notch1*, and *Mill2* (Pan et al. 2017). Increased production of inflammatory mediators (IL-6) and bacterial translocation across gut mucosa may enhance leukemogenesis in *Tet2*^{-/-} mice (Zhang et al. 2015; Meisel et al. 2018). Recent studies also indicate aberrant expression of diverse metabolic pathways, such as BCAT1, can decrease TET function by inhibiting α -ketoglutarate (Raffel et al. 2017). Supplementation of vitamin C (ascorbate), a co-factor of α -ketoglutarate-dependent dioxygenases like TET enzymes, has recently been shown to enhance residual *TET2* function in the heterozygous mutant state (Agathocleous et al. 2017; Cimmino et al. 2017). A clinical trial evaluating the safety and efficacy of high-dose vitamin C for *TET2*-mutated myeloid neoplasms is ongoing. Despite our mechanistic understanding of the function of TET proteins, how loss-of-function mutations in *TET2* initiates and maintains leukemogenesis is largely unknown.

IDH1 and IDH2 Mutations

High-throughput sequencing of AML patient samples identified neomorphic mutations in cytosolic (*IDH1* at R132) or mitochondrial (*IDH2* at R140Q or R172) isoforms of isocitrate dehydrogenase, which are critical for the oxidative

A. Kishtagari and R.L. Levine

carboxylation of isocitrate to α -ketoglutarate. These mutations contribute to leukemogenesis through the production of the oncometabolite R enantiomer of 2-hydroxyglutamate (R-2-HG), which, in turn, impedes epigenetic regulation via competitive inhibition of dioxygenases, including the TET family of enzymes and histone lysine demethylases (KDM), by competing with α -ketoglutarate (Mardis et al. 2009; Figueroa et al. 2010; Xu et al. 2011). This leads to a repressive chromatin landscape via DNA and histone hypermethylation, and consequent block in cellular differentiation (Lu et al. 2012; Losman et al. 2013). Mutations in *IDH1* and *IDH2* are present in ~5%–10% and ~15%–20% of patients with newly diagnosed AML, respectively. Mutations in *IDH1* and *IDH2* mutually exclusive and occur as early clonal events (Papaemmanuil et al. 2016). *IDH1* and *IDH2* mutations are more frequently observed in patients with trisomy 8 and normal karyotype AML (Medeiros et al. 2017). First-in-class, oral, selective inhibitors of mutant *IDH1* (ivosidenib) and *IDH2* (enasidenib) are U.S. Federal Drug Administration (FDA)-approved for the management of *IDH1/2*-mutant adults with refractory or relapsed AML (Stein et al. 2017; DiNardo et al. 2018). Several mechanisms of resistance to enasidenib (*IDH2* inhibitor) leading to late relapse have already been described, including acquisition of *IDH1*-mutated subclones (Quek et al. 2018) or development of additional noncatalytic second-site mutations of *IDH2* (Intlekofer et al. 2018).

ASXL1

Additional sex combs-like 1 (*ASXL1*), located at 20q11.21, is one of the three mammalian homologs of the *Drosophila* additional sex combs (*Asx*) gene. *Asx* functions as an epigenetic regulator that was originally identified as an enhancer of *Trithorax* (*TrxG*) and Polycomb group (*PcG*) genes (Micol and Abdel-Wahab 2016). *ASXL1* interacts with BRCA1-associated protein 1 (BAP1) to form the Polycomb repressive deubiquitinase (PR-DUB) complex, which removes monoubiquitin from histone H2A at lysine 119 (H2AK119Ub) (a repressive mark)

to activate genes targeted by PRC1 (Scheuermann et al. 2010). *ASXL1* also interacts with the PRC2 complex to mediate histone H3 lysine 27 (H3K27) trimethylation (Abdel-Wahab et al. 2012; Inoue et al. 2013). It is yet to be determined if the observed changes in H3K27 methylation with *ASXL1* loss are caused by direct interaction between *ASXL1* and PRC2 or are secondary to depletion of H2AK119Ub and subsequent failure of PRC2 recruitment (Bala-subramani et al. 2015). Somatic mutations in *ASXL1* are frequently detected in clonal hematopoiesis and various myeloid malignancies, suggesting they are the initial events in the leukemic transformation. *ASXL1* mutations occur in ~10%–20% of patients with AML and are enriched in those with underlying myelodysplasia (Metzeler et al. 2011). *ASXL1* mutations coexist with those mutations in the splicing factor (*SRSF2*, *U2AF1*), signal transduction (*NRAS*, *JAK2*, *NF1*), and transcription factor (*RUNX1*) (Papaemmanuil et al. 2016). *ASXL1* mutations are more common in older patients with AML and confer a poor prognosis (Schnittger et al. 2013). A mechanistic understanding of how *ASXL1* mutations in myeloid malignancies drive leukemogenesis remains to be fully elucidated.

BCOR and BCORL1

BCL6 corepressor (*BCOR*) and its closely related homolog, BCL6 corepressor like 1 (*BCORL1*), are located on the X-chromosome and are targeted by somatic mutations in AML, MDS, chronic myelomonocytic leukemia (CMML), and aplastic anemia (Damm et al. 2013; Yoshizato et al. 2015; Papaemmanuil et al. 2016). *BCOR/BCORL1* function as components of PRC1.1, a noncanonical PRC1 complex. PRC1.1 is responsible for the deposition of monoubiquitination on lysine 119 of histone 2A (H2AK119ub1) and mediates transcriptional repression by recruitment of PRC2 and subsequent deposition of H3K27me3 mark (Blackledge et al. 2014; Di Carlo et al. 2019). In mice, loss of *Bcor* function results in enhanced self-renewal of myeloid progenitor cells and cooperates with *Kras*^{G12D} to initiate leukemia in vivo (Kelly et al. 2019).

Loss-of-function somatic mutations in *BCOR* and *BCORL1* have been reported in ~1%–4% of AML, and carry an unfavorable prognosis (Grossmann et al. 2011; Damm et al. 2013).

Splicing Factor Mutations (*SF3B1*, *SRSF2*, *U2AF1*, *ZRSR2*, *U2AF2*, *PRPF8*, and *LUC7L2*)

Almost all human genes are subject to alternative pre-mRNA splicing, a primary source of diversity in messenger RNA species and protein isoforms, which is orchestrated by the macromolecular spliceosome complex (Fig. 1) (Lee and Rio 2015). Recent advances in high-throughput sequencing technologies have unexpectedly identified somatic mutations in the core spliceosomal proteins and associated RNA splicing factors in MDS, CMML, and AML (Anczuków and Krainer 2016). This includes mutations in *SF3B1* (splicing factor 3b subunit 1), *SRSF2* (serine/arginine rich splicing factor 2), *U2AF1* (U2 small nuclear RNA auxiliary factor 1), and *ZRSR2* (zinc finger RNA binding motif and serine/arginine rich 2). These mutations are mutually exclusive, heterozygous point mutations, suggesting synthetic lethal interactions if they were to co-occur (Yoshida et al. 2011; Lee et al. 2018). Splicing machinery mutations are most common in AML arising out of antecedent myeloid malignancy (secondary AML) (Linsley et al. 2015). It has been shown that mutations in *SRSF2* cooperate with *IDH2*, with resultant synergy between RNA splicing and epigenetic regulation in promoting leukemogenesis (Yoshimi et al. 2019). Several groups have developed therapeutic strategies targeting the core spliceosome (*SF3B1* inhibitors), splicing regulatory factors (*PRMT5* inhibitor, *RBM39* degraders, and *SRPKs/CLKs* inhibitors), and modulating pathologic splicing events (oligonucleotide-based approaches) (Lee et al. 2016; Obeng et al. 2016; Seiler et al. 2018; Fong et al. 2019; Wang et al. 2019).

Cohesin Complex

The cohesin complex is a tripartite ring including three structural proteins, *SMC1A*, *SMC3*,

Somatic Mutations in Acute Myeloid Leukemia

and *RAD21*, bound to either *STAG1* or *STAG2* (Fig. 2). This complex has many functions, most notably an essential role in aligning and stabilizing sister chromatids during metaphase (Nasmyth and Haering 2009). Cohesin complex also plays a critical role in the regulation of gene expression by cooperating with the DNA-binding protein CTCF in maintaining topologically associated domains (TADs) and regulating three-dimensional (3D) genome organization (Wendt et al. 2008). Heterozygous loss-of-function mutations in the cohesin complex occur in 12%–20% of AML and MDS patients and are more prevalent in high-risk MDS and secondary AML (Kon et al. 2013; Thota et al. 2014). These mutations are not associated with aneuploidy. Several studies have reported that the haploinsufficiency induced by the cohesin mutations enhances HSPC self-renewal and block hematopoietic differentiation (Mazumdar et al. 2015; Mullenders et al. 2015; Viny et al. 2015). *STAG2*, which is X-linked, is the most commonly mutated cohesin gene across many hematopoietic and solid tumors. A recent study demonstrated a specific role for *STAG2* (not by *STAG1*) in balancing self-renewal and differentiation in HSPCs (Viny et al. 2019).

Mutations in Tumor-Suppressor Genes: *TP53*, *WT1*, *PHF6*

TP53

The tumor-suppressor gene *TP53* is the most frequently mutated gene in human cancer and has many cellular functions, which are discussed in detail elsewhere (Fig. 1) (Kasthuber and Lowe 2017). Mutations in *TP53* occur in 5%–20% of patients with newly diagnosed AML. There is an increased frequency of *TP53* mutations in AML patients with therapy-related myeloid neoplasms, complex karyotype, monosomal karyotype, and relapsed disease. One possible postulation for *TP53* pathogenesis in therapy-related myeloid neoplasms is that *TP53* mutations are acquired in some HSPCs during normal aging (clonal hematopoiesis) and are then selected for upon exposure to cytotoxic chemotherapy, ultimately resulting in

A. Kishtagari and R.L. Levine

the expansion of HSPCs with *TP53* mutations (Wong et al. 2015). The majority of *TP53* mutations are missense mutations within the central DNA-binding domain, leading to loss of function. The resulting loss of function of p53 favors genomic instability and impaired apoptosis, which can contribute to resistance to chemotherapy. AML patients with *TP53* mutations have been associated with an inadequate response to conventional cytotoxic therapies and inferior overall survival rates (median of 5–9 mo) (Stengel et al. 2017). Historically, the efforts to target p53 therapeutically have been challenging. The landscape has been changing with the development of novel therapeutic strategies targeting the *TP53* pathway. One such promising approach is APR-246, a methylated derivative of PRIMA-1 that is converted to Michael acceptor methylene quinuclidinone (MQ), which can bind covalently to mutant p53 and restore wild-type transcriptional properties to the protein (Lambert et al. 2009). There is an ongoing multicenter phase III randomized study comparing the combination of APR-246 with azacytidine compared with azacytidine alone in *TP53*-mutant MDS and AML (NCT03745716).

WT1

Wilm's tumor 1 (*WT1*) encodes for a transcription factor that regulates cell proliferation, differentiation, apoptosis, and organ development (Call et al. 1990). Somatic loss-of-function mutations target *WT1* in 6%–15% of AML patients. Also, overexpression of wild-type *WT1* protein has been observed in AML, further challenging the mechanistic understanding of how *WT1* contributes to leukemogenesis (Rampal and Figueroa 2016). Heterozygous loss of *Wt1* enhances HSCs self-renewal and cooperates with *Flt3-ITD* to induce leukemic transformation (Pronier et al. 2018). Loss-of-function mutations in *WT1* led to marked promoter hypermethylation pattern and reduction in 5hmC levels similar to AML patients with mutations in *TET2* and *IDH1/2* (Rampal et al. 2014). *WT1* protein has also been shown to bind directly to *TET2* and recruit *TET2* to specific genomic sites to activate the gene expression of *WT1*-target

genes (Wang et al. 2015). Mutations in *WT1* are associated with younger age, often co-occur with *FLT3-ITD*, and confer adverse prognostic significance in AML with increased risk of primary refractory disease (Hou et al. 2010).

PHF6

Somatic loss-of-function mutations in the X-chromosome-linked tumor suppressor plant homeodomain factor 6 (*PHF6*) are present in ~3% of AML patients (Van Vlierberghe et al. 2011). *PHF6* is involved in chromatin-mediated transcriptional regulation by binding to the nucleosome remodeling and deacetylase (NuRD) chromatin remodeling complex (Todd and Picketts 2012; Liu et al. 2015). The presence of *PHF6* mutations in preleukemic clonal hematopoiesis supports its role in leukemia initiation and HSC self-renewal (Yoshizato et al. 2015; Abelson et al. 2018). In line with this notion, in a mouse model of T-cell acute lymphoblastic anemia (T-ALL), loss of *Phf6* enhances tumor initiation and transformation and leads to increased self-renewal in mouse HSCs (Wendorff et al. 2019). Mutations in *PHF6* are associated with *RUNX1* mutations and with reduced overall survival in AML patients (Patel et al. 2012).

CONCLUDING REMARKS

Mutational studies involving large cohorts of patients with AML have identified a complex mutational landscape, including patterns of significant co-occurrence and mutual exclusivity of mutations. The mechanistic interactions between different mutations have only begun to be elucidated, and we expect these insights can be used to better inform our understanding of molecular pathogenesis of the disease, improve risk stratification, and identify novel therapeutic vulnerabilities. In specific cases, such as co-occurring *FLT3/TET2* (Shih et al. 2015) and *SRSF2/IDH2* (Yoshimi et al. 2019) mutations, mechanisms of oncogenic synergy have been elucidated; however, the basis for cooperativity between AML disease alleles requires greater investigation. Moreover, bulk sequencing studies

have given us a glimpse into the temporal and spatial sequence of somatic mutations that drive leukemogenesis, from early mutations, which induce clonal hematopoiesis (*DNMT3A*, *TET2*, *ASXL1*), to disease-defining mutations present in all cells at transformation (*NPM1*), to mutations that can occur in specific subclones and further enhance the fitness of AML cells (*FLT3*, *RAS*). However, the roles of all of the different mutations in this complex sequence of genetic events that contribute to AML are not known, and many AML patients are a complex amalgam of multiple branching clones such that the AML “ecosystem” has not been fully characterized. Moreover, the observations that relapse can come from rarer AML subclones (Ding et al. 2012; Shlush et al. 2017) suggest that a better understanding of the role of specific mutations in AML heterogeneity and drug response is urgently needed.

In addition to novel mechanistic insights, we need to ensure that this knowledge is translated into improved outcomes for our patients. Most AML patients do not present with mutations for which we have a molecularly targeted therapy, and targeted therapy has limited curative potential when given as monotherapy in AML. We need to identify additional mechanisms of leukemic transformation induced by genetic/epigenetic events in AML, such that we can uncover novel therapeutic targets that lead to genotype-based combination therapies for AML patients. It is our hope that the ever-evolving understanding of AML pathogenesis will continue to lead to novel therapies for our patients.

ACKNOWLEDGMENTS

This work was supported by National Cancer Institute R35 CA197594-01A1 (R.L.L.), National Cancer Institute R01 CA216421 (R.L.L.), National Cancer Institute PS-OC U54 CA143869-05 (R.L.L.), and Leukemia & Lymphoma Society Translational Research Foundation 6499-17 (R.L.L.). This work was supported in part by Memorial Sloan Kettering Cancer Center Core Grant P30 CA008748. A.K. is supported by the VeloSano Catalyst Award.

Somatic Mutations in Acute Myeloid Leukemia

REFERENCES

- Abdel-Wahab O, Mullally A, Hedvat C, Garcia-Manero G, Patel J, Wadleigh M, Malinge S, Yao J, Kilpivaara O, Bhat R, et al. 2009. Genetic characterization of TET1, TET2, and TET3 alterations in myeloid malignancies. *Blood* **114**: 144–147. doi:10.1182/blood-2009-03-210039
- Abdel-Wahab O, Adli M, LaFave LM, Gao J, Hricik T, Shih AH, Pandey S, Patel JP, Chung YR, Koche R, et al. 2012. *ASXL1* mutations promote myeloid transformation through loss of PRC2-mediated gene repression. *Cancer Cell* **22**: 180–193. doi:10.1016/j.ccr.2012.06.032
- Abelson S, Collord G, Ng SWK, Weissbrod O, Mendelson Cohen N, Niemeyer E, Barda N, Zuzarte PC, Heisler L, Sundaravadanam Y, et al. 2018. Prediction of acute myeloid leukaemia risk in healthy individuals. *Nature* **559**: 400–404. doi:10.1038/s41586-018-0317-6
- Agathocleous M, Meacham CE, Burgess RJ, Piskounova E, Zhao Z, Crane GM, Cowin BL, Bruner E, Murphy MM, Chen W, et al. 2017. Ascorbate regulates haematopoietic stem cell function and leukaemogenesis. *Nature* **549**: 476–481. doi:10.1038/nature23876
- Amatangelo MD, Quek L, Shih A, Stein EM, Roshal M, David MD, Marteyn B, Farnoud NR, de Botton S, Bernard OA, et al. 2017. Enasidenib induces acute myeloid leukemia cell differentiation to promote clinical response. *Blood* **130**: 732–741. doi:10.1182/blood-2017-04-779447
- Anczuków O, Krainer AR. 2016. Splicing-factor alterations in cancers. *RNA* **22**: 1285–1301. doi:10.1261/rna.057919.116
- Arber DA, Orazi A, Hasserjian R, Thiele J, Borowitz MJ, Le Beau MM, Bloomfield CD, Cazzola M, Vardiman JW. 2016. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood* **127**: 2391–2405. doi:10.1182/blood-2016-03-643544
- Armstrong SA, Staunton JE, Silverman LB, Pieters R, den Boer ML, Minden MD, Sallan SE, Lander ES, Golub TR, Korsmeyer SJ. 2002. MLL translocations specify a distinct gene expression profile that distinguishes a unique leukemia. *Nat Genet* **30**: 41–47. doi:10.1038/ng765
- Avellino R, Delwel R. 2017. Expression and regulation of C/EBPα in normal myelopoiesis and in malignant transformation. *Blood* **129**: 2083–2091. doi:10.1182/blood-2016-09-687822
- Bacher U, Haferlach T, Schoch C, Kern W, Schnittger S. 2006. Implications of *NRAS* mutations in AML: A study of 2502 patients. *Blood* **107**: 3847–3853. doi:10.1182/blood-2005-08-3522
- Balasubramani A, Larjo A, Bassein JA, Chang X, Hastie RB, Togher SM, Lähdesmäki H, Rao A. 2015. Cancer-associated *ASXL1* mutations may act as gain-of-function mutations of the *ASXL1*-BAP1 complex. *Nat Commun* **6**: 7307. doi:10.1038/ncomms8307
- Ball BJ, Hsu M, Devlin SM, Famulare C, Cai SF, Dunbar A, Epstein-Peterson ZD, Menghrajani K, Glass JL, Taylor J, et al. 2019. RAS mutations are independently associated with decreased overall survival and event-free survival in patients with AML receiving induction chemotherapy. *Blood* **134**: 18. doi:10.1182/blood-2019-125319
- Basecke J, Whelan JT, Griesinger F, Bertrand FE. 2006. The MLL partial tandem duplication in acute myeloid leukemia.

A. Kishtagari and R.L. Levine

- mia. *Br J Haematol* **135**: 438–449. doi:10.1111/j.1365-2141.2006.06301.x
- Bezerra MF, Lima AS, Piqué-Borràs MR, Silveira DR, Coelho-Silva JL, Pereira-Martins DA, Weinhäuser I, Franca-Neto PL, Quek L, Corby A, et al. 2020. Co-occurrence of DNMT3A, NPM1, FLT3 mutations identifies a subset of acute myeloid leukemia with adverse prognosis. *Blood* **135**: 870–875. doi:10.1182/blood.2019003339
- Blackledge NP, Farcas AM, Kondo T, King HW, McGouran JF, Hanssen LL, Ito S, Cooper S, Kondo K, Koseki Y, et al. 2014. Variant PRC1 complex-dependent H2A ubiquitylation drives PRC2 recruitment and polycomb domain formation. *Cell* **157**: 1445–1459. doi:10.1016/j.cell.2014.05.004
- Bloomfield CD, Lawrence D, Byrd JC, Carroll A, Pettenati MJ, Tantravahi R, Patil SR, Davey FR, Berg DT, Schiffer CA, et al. 1998. Frequency of prolonged remission duration after high-dose cytarabine intensification in acute myeloid leukemia varies by cytogenetic subtype. *Cancer Res* **58**: 4173–4179.
- Borkin D, He S, Miao H, Kempinska K, Pollock J, Chase J, Purohit T, Malik B, Zhao T, Wang J, et al. 2015. Pharmacologic inhibition of the menin-MLL interaction blocks progression of MLL leukemia in vivo. *Cancer Cell* **27**: 589–602. doi:10.1016/j.ccr.2015.02.016
- Brunetti L, Gundry MC, Sorcini D, Guzman AG, Huang YH, Ramabadran R, Gionfriddo I, Mezzasoma F, Milano F, Nabet B, et al. 2018. Mutant NPM1 maintains the leukemic state through HOX expression. *Cancer Cell* **34**: 499–512.e9. doi:10.1016/j.ccr.2018.08.005
- Busque L, Patel JP, Figueroa ME, Vasanthakumar A, Provost S, Hamilou Z, Mollica L, Li J, Viale A, Heguy A, et al. 2012. Recurrent somatic *TET2* mutations in normal elderly individuals with clonal hematopoiesis. *Nat Genet* **44**: 1179–1181. doi:10.1038/ng.2413
- Byrd JC, Ruppert AS, Mrózek K, Carroll AJ, Edwards CG, Arthur DC, Pettenati MJ, Stamborg J, Koduru PR, Moore JO, et al. 2004. Repetitive cycles of high-dose cytarabine benefit patients with acute myeloid leukemia and inv(16)(p13q22) or t(16;16)(p13;q22): results from CALGB 8461. *J Clin Oncol* **22**: 1087–1094. doi:10.1200/JCO.2004.07.012
- Call KM, Glaser T, Ito CY, Buckler AJ, Pelletier J, Haber DA, Rose EA, Kral A, Yeger H, Lewis WH, et al. 1990. Isolation and characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus. *Cell* **60**: 509–520. doi:10.1016/0092-8674(90)90601-A
- Cancer Genome Atlas Research Network, Ley TJ, Miller C, Ding L, Raphael BJ, Mungall AJ, Robertson A, Hoadley K, Triche TJ, Laird PW, et al. 2013. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *New Engl J Med* **368**: 2059–2074. doi:10.1056/NEJMoa1301689
- Challen GA, Sun D, Jeong M, Luo M, Jelinek J, Berg JS, Bock C, Vasanthakumar A, Gu H, Xi Y, et al. 2011. Dnmt3a is essential for hematopoietic stem cell differentiation. *Nat Genet* **44**: 23–31. doi:10.1038/ng.1009
- Chen TH, Kambal A, Krysiak K, Walshauser MA, Raju G, Tibbitts JF, Walter MJ. 2011. Knockdown of Hspa9, a del(5q31.2) gene, results in a decrease in hematopoietic progenitors in mice. *Blood* **117**: 1530–1539. doi:10.1182/blood-2010-06-293167
- Chen C, Liu Y, Rappaport AR, Kitzing T, Schultz N, Zhao Z, Shroff AS, Dickins RA, Vakoc CR, Bradner JE, et al. 2014. MLL3 is a haploinsufficient 7q tumor suppressor in acute myeloid leukemia. *Cancer Cell* **25**: 652–665. doi:10.1016/j.ccr.2014.03.016
- Chen L, Chen W, Mysliwski M, Serio J, Ropa J, Abulwerdi FA, Chan RJ, Patel JP, Tallman MS, Paietta E, et al. 2015. Mutated *Ptfn11* alters leukemic stem cell frequency and reduces the sensitivity of acute myeloid leukemia cells to Mcl1 inhibition. *Leukemia* **29**: 1290–1300. doi:10.1038/leu.2015.18
- Cichowski K, Jacks T. 2001. NF1 tumor suppressor gene function: Narrowing the GAP. *Cell* **104**: 593–604. doi:10.1016/S0092-8674(01)00245-8
- Cimmino L, Dolgalev I, Wang Y, Yoshimi A, Martin GH, Wang J, Ng V, Xia B, Witkowski MT, Mitchell-Flack M, et al. 2017. Restoration of *TET2* function blocks aberrant self-renewal and leukemia progression. *Cell* **170**: 1079–1095.e20. doi:10.1016/j.cell.2017.07.032
- Crispino JD, Horwitz MS. 2017. GATA factor mutations in hematologic disease. *Blood* **129**: 2103–2110. doi:10.1182/blood-2016-09-687889
- Crute BE, Lewis AF, Wu Z, Bushweller JH, Speck NA. 1996. Biochemical and biophysical properties of the core-binding factor $\alpha 2$ (AML1) DNA-binding domain. *J Biol Chem* **271**: 26251–26260. doi:10.1074/jbc.271.42.26251
- Damm F, Chesnais V, Nagata Y, Yoshida K, Scourzic L, Okuno Y, Itzykson R, Sanada M, Shiraiishi Y, Gelsi-Boyer V, et al. 2013. *BCOR* and *BCORL1* mutations in myelodysplastic syndromes and related disorders. *Blood* **122**: 3169–3177. doi:10.1182/blood-2012-11-469619
- Delhommeau F, Dupont S, Della Valle V, James C, Trannoy S, Massé A, Kosmider O, Le Couedic JP, Robert F, Alberdi A, et al. 2009. Mutation in *TET2* in myeloid cancers. *New Engl J Med* **360**: 2289–2301. doi:10.1056/NEJMoa0810069
- Desai P, Mencia-Trinchant N, Savenkov O, Simon MS, Cheang G, Lee S, Samuel M, Ritchie EK, Guzman ML, Ballman KV, et al. 2018. Somatic mutations precede acute myeloid leukemia years before diagnosis. *Nat Med* **24**: 1015–1023. doi:10.1038/s41591-018-0081-z
- De Weer A, Speleman F, Cauwelier B, Van Roy N, Yigit N, Verhasselt B, De Moerloose B, Benoit Y, Noens L, Sellsag D, et al. 2008. *EVII* overexpression in t(3;17) positive myeloid malignancies results from juxtaposition of *EVII* to the *MSI2* locus at 17q22. *Haematologica* **93**: 1903–1907. doi:10.3324/haematol.13192
- Díaz-Beyá M, Labopin M, Maertens J, Alijurf M, Passweg J, Dietrich B, Schouten H, Socié G, Schaap N, Schwerdtfeger R, et al. 2020. Allogeneic stem cell transplantation in AML with t(6;9)(p23;q34);*DEK-NUP214* shows a favourable outcome when performed in first complete remission. *Br J Haematol* doi: 10.1111/bjh.16433
- Di Carlo V, Mocavini I, Di Croce L. 2019. Polycomb complexes in normal and malignant hematopoiesis. *J Cell Biol* **218**: 55–69. doi:10.1083/jcb.201808028
- Dillon R, Hills R, Freeman S, Potter N, Jovanovic J, Ivey A, Kanda AS, Rungllall M, Foot N, Valganon M, et al. 2020. Molecular MRD status and outcome after transplantation in *NPM1*-mutated AML. *Blood* **135**: 680–688. doi:10.1182/blood.2019002959

- DiNardo CD, Stein EM, de Botton S, Roboz GJ, Altman JK, Mims AS, Swords R, Collins RH, Mannis GN, Pollyea DA, et al. 2018. Durable remissions with ivosidenib in *IDH1*-mutated relapsed or refractory AML. *New Engl J Med* **378**: 2386–2398. doi:10.1056/NEJMoa1716984
- DiNardo CD, Tiong IS, Quaglieri A, MacRaid S, Lohavi S, Brown FC, Thijssen R, Pomilio G, Ivey A, Salmon JM, et al. 2020. Molecular patterns of response and treatment failure after frontline venetoclax combinations in older patients with AML. *Blood* **135**: 791–803. doi:10.1182/blood.2019003988
- Ding L, Ley TJ, Larson DE, Miller CA, Koboldt DC, Welch JS, Ritchey JK, Young MA, Lamprecht T, McLellan MD, et al. 2012. Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* **481**: 506–510. doi:10.1038/nature10738
- Döhner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, Büchner T, Dombret H, Ebert BL, Fenaux P, Larson RA, et al. 2017. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood* **129**: 424–447. doi:10.1182/blood-2016-08-733196
- Ebert BL, Pretz J, Bosco J, Chang CY, Tamayo P, Galili N, Raza A, Root DE, Attar E, Ellis SR, et al. 2008. Identification of RPS14 as a 5q- syndrome gene by RNA interference screen. *Nature* **451**: 335–339. doi:10.1038/nature06494
- Eisfeld AK, Kohlschmidt J, Mrózek K, Mims A, Walker CJ, Blachly JS, Nicolet D, Orwick S, Maharry SE, Carroll AJ, et al. 2018. *NF1* mutations are recurrent in adult acute myeloid leukemia and confer poor outcome. *Leukemia* **32**: 2536–2545. doi:10.1038/s41375-018-0147-4
- Falini B, Mecucci C, Tiacci E, Alcalay M, Rosati R, Pasqualucci L, La Starza R, Diverio D, Colombo E, Santucci A, et al. 2005. Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *New Engl J Med* **352**: 254–266. doi:10.1056/NEJMoa041974
- Fasan A, Haferlach C, Alpermann T, Jeromin S, Grossmann V, Eder C, Weissmann S, Dicker F, Kohlmann A, Schindela S, et al. 2014. The role of different genetic subtypes of *CEBPA* mutated AML. *Leukemia* **28**: 794–803. doi:10.1038/leu.2013.273
- Figuerola ME, Abdel-Wahab O, Lu C, Ward PS, Patel J, Shih A, Li Y, Bhagwat N, Vasanthakumar A, Fernandez HF, et al. 2010. Leukemic *IDH1* and *IDH2* mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell* **18**: 553–567. doi:10.1016/j.ccr.2010.11.015
- Fong JY, Pignata L, Goy PA, Kawabata KC, Lee SC, Koh CM, Musiani D, Massignani E, Kotini AG, Penson A, et al. 2019. Therapeutic targeting of RNA splicing catalysis through inhibition of protein arginine methylation. *Cancer Cell* **36**: 194–209.e9. doi:10.1016/j.ccell.2019.07.003
- Fröhling S, Schlenk RF, Stolze I, Bihlmayr J, Benner A, Kreitmeier S, Tobis K, Döhner H, Döhner K. 2004. *CEBPA* mutations in younger adults with acute myeloid leukemia and normal cytogenetics: Prognostic relevance and analysis of cooperating mutations. *J Clin Oncol* **22**: 624–633. doi:10.1200/JCO.2004.06.060
- Gaidzik VI, Teleanu V, Papaemmanuil E, Weber D, Paschka P, Hahn J, Wallrabenstein T, Kolbinger B, Köhne CH, Horst HA, et al. 2016. *RUNX1* mutations in acute myeloid leukemia are associated with distinct clinico-pathologic and genetic features. *Leukemia* **30**: 2160–2168. doi:10.1038/leu.2016.126
- Genovese G, Kähler AK, Handsaker RE, Lindberg J, Rose SA, Bakhoum SF, Chambert K, Mick E, Neale BM, Fromer M, et al. 2014. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *New Engl J Med* **371**: 2477–2487. doi:10.1056/NEJMoa1409405
- Greif PA, Dufour A, Konstandin NP, Ksienzyk B, Zellmeier E, Tizazu B, Sturm J, Benthaus T, Herold T, Yaghmaie M, et al. 2012. *GATA2* zinc finger 1 mutations associated with biallelic *CEBPA* mutations define a unique genetic entity of acute myeloid leukemia. *Blood* **120**: 395–403. doi:10.1182/blood-2012-01-403220
- Grembecka J, He S, Shi A, Purohit T, Muntean AG, Sorenson RJ, Showalter HD, Murai MJ, Belcher AM, Hartley T, et al. 2012. Menin-MLL inhibitors reverse oncogenic activity of MLL fusion proteins in leukemia. *Nat Chem Biol* **8**: 277–284. doi:10.1038/nchembio.773
- Gröschel S, Sanders MA, Hoogenboezem R, de Wit E, Bouwman BAM, Erpelinck C, van der Velden VHJ, Havermans M, Avellino R, van Lom K, et al. 2014. A single oncogenic enhancer rearrangement causes concomitant *EVII* and *GATA2* deregulation in leukemia. *Cell* **157**: 369–381. doi:10.1016/j.cell.2014.02.019
- Gröschel S, Sanders MA, Hoogenboezem R, Zeilemaker A, Havermans M, Erpelinck C, Bindels EM, Beverloo HB, Döhner H, Löwenberg B, et al. 2015. Mutational spectrum of myeloid malignancies with *inv(3)(t(3;3))* reveals a predominant involvement of RAS/RTK signaling pathways. *Blood* **125**: 133–139. doi:10.1182/blood-2014-07-591461
- Grossmann V, Tiacci E, Holmes AB, Kohlmann A, Martelli MP, Kern W, Spanhol-Rosseto A, Klein HU, Dugas M, Schindela S, et al. 2011. Whole-exome sequencing identifies somatic mutations of *BCOR* in acute myeloid leukemia with normal karyotype. *Blood* **118**: 6153–6163. doi:10.1182/blood-2011-07-365320
- Gu X, Ebrahim Q, Mahfouz RZ, Hasipek M, Enane F, Radivoyevitch T, Rapin N, Przychodzen B, Hu Z, Balusu R, et al. 2018. Leukemogenic nucleophosmin mutation disrupts the transcription factor hub that regulates granulomonocytic fates. *J Clin Invest* **128**: 4260–4279. doi:10.1172/JCI97117
- Guryanova OA, Shank K, Spitzer B, Luciani L, Koche RP, Garrett-Bakelman FE, Ganzel C, Durham BH, Mohanty A, Hoermann G, et al. 2016. *DNMT3A* mutations promote anthracycline resistance in acute myeloid leukemia via impaired nucleosome remodeling. *Nat Med* **22**: 1488–1495. doi:10.1038/nm.4210
- Hahn CN, Chong CE, Carmichael CL, Wilkins EJ, Brautigan PJ, Li XC, Babic M, Lin M, Carmagnac A, Lee YK, et al. 2011. Heritable *GATA2* mutations associated with familial myelodysplastic syndrome and acute myeloid leukemia. *Nat Genet* **43**: 1012–1017. doi:10.1038/ng.913
- Hatlen MA, Arora K, Vacic V, Grabowska EA, Liao W, Riley-Gillis B, Oschwald DM, Wang L, Joergens JE, Shih AH, et al. 2016. Integrative genetic analysis of mouse and human AML identifies cooperating disease alleles. *J Exp Med* **213**: 25–34. doi:10.1084/jem.20150524
- He YF, Li BZ, Li Z, Liu P, Wang Y, Tang Q, Ding J, Jia Y, Chen Z, Li L, et al. 2011. Tet-mediated formation of 5-carbox-

A. Kishtagari and R.L. Levine

- ylcytosine and its excision by TDG in mammalian DNA. *Science* **333**: 1303–1307. doi:10.1126/science.1210944
- Hinai A, Pratcorona M, Grob T, Kavelaars FG, Bussaglia E, Sanders MA, Nomdedeu J, Valk PJM. 2019. The landscape of *KMT2A*-PTD AML: Concurrent mutations, gene expression signatures, and clinical outcome. *Hemisphere* **3**: e181. doi:10.1097/HS9.0000000000000181
- Hirsch P, Zhang Y, Tang R, Joulin V, Boutroux H, Pronier E, Moatti H, Flandrin P, Marzac C, Bories D, et al. 2016. Genetic hierarchy and temporal variegation in the clonal history of acute myeloid leukaemia. *Nat Commun* **7**: 12475. doi:10.1038/ncomms12475
- Holz-Schietinger C, Matje DM, Reich NO. 2012. Mutations in DNA methyltransferase (DNMT3A) observed in acute myeloid leukemia patients disrupt processive methylation. *J Biol Chem* **287**: 30941–30951. doi:10.1074/jbc.M112.366625
- Hosono N, Makishima H, Jerez A, Yoshida K, Przyschodzen B, McMahon S, Shiraishi Y, Chiba K, Tanaka H, Miyano S, et al. 2014. Recurrent genetic defects on chromosome 7q in myeloid neoplasms. *Leukemia* **28**: 1348–1351. doi:10.1038/leu.2014.25
- Hou HA, Huang TC, Lin LI, Liu CY, Chen CY, Chou WC, Tang JL, Tseng MH, Huang CF, Chiang YC, et al. 2010. *WT1* mutation in 470 adult patients with acute myeloid leukemia: Stability during disease evolution and implication of its incorporation into a survival scoring system. *Blood* **115**: 5222–5231. doi:10.1182/blood-2009-12-259390
- Inoue D, Kitaura J, Togami K, Nishimura K, Enomoto Y, Uchida T, Kagiya Y, Kawabata KC, Nakahara F, Izawa K, et al. 2013. Myelodysplastic syndromes are induced by histone methylation-altering *ASXL1* mutations. *J Clin Invest* **123**: 4627–4640. doi:10.1172/JCI70739
- Intlekofer AM, Shih AH, Wang B, Nazir A, Rustenburg AS, Albanese SK, Patel M, Famulare C, Correa FM, Takemoto N, et al. 2018. Acquired resistance to IDH inhibition through *trans* or *cis* dimer-interface mutations. *Nature* **559**: 125–129. doi:10.1038/s41586-018-0251-7
- Ito S, Shen L, Dai Q, Wu SC, Collins LB, Swenberg JA, He C, Zhang Y. 2011. Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* **333**: 1300–1303. doi:10.1126/science.1210597
- Ito Y, Bae SC, Chuang LS. 2015. The RUNX family: Developmental regulators in cancer. *Nat Rev Cancer* **15**: 81–95. doi:10.1038/nrc3877
- Ivey A, Hills RK, Simpson MA, Jovanovic JV, Gilkes A, Grech A, Patel Y, Bhudia N, Farah H, Mason J, et al. 2016. Assessment of minimal residual disease in standard-risk AML. *New Engl J Med* **374**: 422–433. doi:10.1056/NEJMoa1507471
- Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, Lindsley RC, Mermel CH, Burt N, Chavez A, et al. 2014. Age-related clonal hematopoiesis associated with adverse outcomes. *New Engl J Med* **371**: 2488–2498. doi:10.1056/NEJMoa1408617
- Jankowska AM, Szpurka H, Tiu RV, Makishima H, Afable M, Huh J, O'Keefe CL, Ganetzky R, McDevitt MA, Maciejewski JP. 2009. Loss of heterozygosity 4q24 and *TET2* mutations associated with myelodysplastic/myeloproliferative neoplasms. *Blood* **113**: 6403–6410. doi:10.1182/blood-2009-02-205690
- Joslin JM, Fernald AA, Tennant TR, Davis EM, Kogan SC, Anastasi J, Crispino JD, Le Beau MM. 2007. Haploinsufficiency of *EGR1*, a candidate gene in the del(5q), leads to the development of myeloid disorders. *Blood* **110**: 719–726. doi:10.1182/blood-2007-01-068809
- Kastenhuber ER, Lowe SW. 2017. Putting p53 in context. *Cell* **170**: 1062–1078. doi:10.1016/j.cell.2017.08.028
- Kayser S, Hills RK, Lusk MR, Brunner AM, Terré C, Westermann J, Menghrajani K, Shaw C, Baer MR, Elliott MA, et al. 2020. Allogeneic hematopoietic cell transplantation improves outcome of adults with t(6;9) acute myeloid leukemia: Results from an international collaborative study. *Haematologica* **105**: 161–169. doi:10.3324/haematol.2018.208678
- Kelly MJ, So J, Rogers AJ, Gregory G, Li J, Zethoven M, Gearhart MD, Bardwell VJ, Johnstone RW, Vervoort SJ, et al. 2019. Bcor loss perturbs myeloid differentiation and promotes leukaemogenesis. *Nat Commun* **10**: 1347. doi:10.1038/s41467-019-09250-6
- Kirstetter P, Schuster MB, Bereshchenko O, Moore S, Dvinge H, Kurz E, Theilgaard-Mönch K, Månsson R, Pedersen TA, Pabst T, et al. 2008. Modeling of C/EBPα mutant acute myeloid leukemia reveals a common expression signature of committed myeloid leukemia-initiating cells. *Cancer Cell* **13**: 299–310. doi:10.1016/j.ccr.2008.02.008
- Kishtagari A, Levine RL, Viny AD. 2020. Driver mutations in acute myeloid leukemia. *Curr Opin Hematol* **27**: 49–57. doi:10.1097/MOH.0000000000000567
- Klossowski S, Miao H, Kempinska K, Wu T, Purohit T, Kim E, Linhares BM, Chen D, Jih G, Perkey E, et al. 2020. Menin inhibitor MI-3454 induces remission in *MLL1*-rearranged and *NPM1*-mutated models of leukemia. *J Clin Invest* **130**: 981–997. doi:10.1172/JCI129126
- Ko M, Huang Y, Jankowska AM, Pape UJ, Tahiliani M, Bandukwala HS, An J, Lamperti ED, Koh KP, Ganetzky R, et al. 2010. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant *TET2*. *Nature* **468**: 839–843. doi:10.1038/nature09586
- Kon A, Shih LY, Minamino M, Sanada M, Shiraishi Y, Nagata Y, Yoshida K, Okuno Y, Bando M, Nakato R, et al. 2013. Recurrent mutations in multiple components of the cohesin complex in myeloid neoplasms. *Nat Genet* **45**: 1232–1237. doi:10.1038/ng.2731
- Krivtsov AV, Armstrong SA. 2007. *MLL* translocations, histone modifications and leukaemia stem-cell development. *Nat Rev Cancer* **7**: 823–833. doi:10.1038/nrc2253
- Krivtsov AV, Evans K, Gadrey JY, Eschle BK, Hatton C, Uckelmann HJ, Ross KN, Perner F, Olsen SN, Pritchard T, et al. 2019. A Menin-*MLL* inhibitor induces specific chromatin changes and eradicates disease in models of *MLL*-rearranged leukemia. *Cancer Cell* **36**: 660–673.e11. doi:10.1016/j.ccell.2019.11.001
- Kunimoto H, Meydan C, Nazir A, Whitfield J, Shank K, Rapaport F, Maher R, Pronier E, Meyer SC, Garrett-Bakelman FE, et al. 2018. Cooperative epigenetic remodeling by *TET2* loss and *NRAS* mutation drives myeloid transformation and MEK inhibitor sensitivity. *Cancer Cell* **33**: 44–59.e8. doi:10.1016/j.ccell.2017.11.012
- Lambert JM, Gorzov P, Veprintsev DB, Söderqvist M, Segerback D, Bergman J, Fersht AR, Hainaut P, Wiman KG, Bykov VJ. 2009. PRIMA-1 reactivates mutant p53 by co-

- valent binding to the core domain. *Cancer Cell* **15**: 376–388. doi:10.1016/j.ccr.2009.03.003
- Lavallée VP, Kros J, Lemieux S, Boucher G, Gendron P, Pabst C, Boivin I, Marinier A, Guidos CJ, Meloche S, et al. 2016. Chemo-genomic interrogation of *CEBPA* mutated AML reveals recurrent *CSF3R* mutations and subgroup sensitivity to JAK inhibitors. *Blood* **127**: 3054–3061. doi:10.1182/blood-2016-03-705053
- Lee Y, Rio DC. 2015. Mechanisms and regulation of alternative pre-mRNA splicing. *Annu Rev Biochem* **84**: 291–323. doi:10.1146/annurev-biochem-060614-034316
- Lee SC, Dvinge H, Kim E, Cho H, Micol JB, Chung YR, Durham BH, Yoshimi A, Kim YJ, Thomas M, et al. 2016. Modulation of splicing catalysis for therapeutic targeting of leukemia with mutations in genes encoding spliceosomal proteins. *Nat Med* **22**: 672–678. doi:10.1038/nm.4097
- Lee SC, North K, Kim E, Jang E, Obeng E, Lu SX, Liu B, Inoue D, Yoshimi A, Ki M, et al. 2018. Synthetic lethal and convergent biological effects of cancer-associated spliceosomal gene mutations. *Cancer Cell* **34**: 225–241.e8. doi:10.1016/j.ccell.2018.07.003
- Ley TJ, Mardis ER, Ding L, Fulton B, McLellan MD, Chen K, Dooling D, Dunford-Shore BH, McGrath S, Hickenbotham M, et al. 2008. DNA sequencing of a cytogenetically normal acute myeloid leukaemia genome. *Nature* **456**: 66–72. doi:10.1038/nature07485
- Ley TJ, Ding L, Walter MJ, McLellan MD, Lamprecht T, Larson DE, Kandoth C, Payton JE, Baty J, Welch J, et al. 2010. *DNMT3A* mutations in acute myeloid leukemia. *New Engl J Med* **363**: 2424–2433. doi:10.1056/NEJMoa1005133
- Li S, Balmain A, Counter CM. 2018. A model for RAS mutation patterns in cancers: Finding the sweet spot. *Nat Rev Cancer* **18**: 767–777. doi:10.1038/s41568-018-0076-6
- Lindsley RC, Mar BG, Mazzola E, Grauman PV, Shareef S, Allen SL, Pignaux A, Wetzler M, Stuart RK, Erba HP, et al. 2015. Acute myeloid leukemia ontogeny is defined by distinct somatic mutations. *Blood* **125**: 1367–1376. doi:10.1182/blood-2014-11-610543
- Liu TX, Becker MW, Jelinek J, Wu WS, Deng M, Mikhalkovich N, Hsu K, Bloomfield CD, Stone RM, DeAngelo DJ, et al. 2007. Chromosome 5q deletion and epigenetic suppression of the gene encoding α -catenin (CTNNA1) in myeloid cell transformation. *Nat Med* **13**: 78–83. doi:10.1038/nm1512
- Liu Z, Li F, Zhang B, Li S, Wu J, Shi Y. 2015. Structural basis of plant homeodomain finger 6 (PHF6) recognition by the retinoblastoma binding protein 4 (RBBP4) component of the nucleosome remodeling and deacetylase (NuRD) complex. *J Biol Chem* **290**: 6630–6638. doi:10.1074/jbc.M114.610196
- Liu Y, Chen C, Xu Z, Scuoppo C, Rillahan CD, Gao J, Spitzer B, Bosbach B, Kastenhuber ER, Baslan T, et al. 2016. Deletions linked to *TP53* loss drive cancer through p53-independent mechanisms. *Nature* **531**: 471–475. doi:10.1038/nature17157
- Losman JA, Looper RE, Koivunen P, Lee S, Schneider RK, McMahon C, Cowley GS, Root DE, Ebert BL, Kaelin WG Jr. 2013. (R)-2-hydroxyglutarate is sufficient to promote leukemogenesis and its effects are reversible. *Science* **339**: 1621–1625. doi:10.1126/science.1231677
- Lu C, Ward PS, Kapoor GS, Rohle D, Turcan S, Abdel-Wahab O, Edwards CR, Khanin R, Figueroa ME, Melnick A, et al. 2012. IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature* **483**: 474–478. doi:10.1038/nature10860
- Lugthart S, Gröschel S, Beverloo HB, Kayser S, Valk PJ, van Zelderen-Bhola SL, Jan Ossenkoppele G, Vellenga E, van den Berg-de Ruyter E, Schanz U, et al. 2010. Clinical, molecular, and prognostic significance of WHO type inv(3)(q21q26.2)/t(3;3)(q21;q26.2) and various other 3q abnormalities in acute myeloid leukemia. *J Clin Oncol* **28**: 3890–3898. doi:10.1200/JCO.2010.29.2771
- Makishima H, Yoshizato T, Yoshida K, Sekeres MA, Radivoyevitch T, Suzuki H, Przyschodzen B, Nagata Y, Megendorfer M, Sanada M, et al. 2017. Dynamics of clonal evolution in myelodysplastic syndromes. *Nat Genet* **49**: 204–212. doi:10.1038/ng.3742
- Malaise M, Steinbach D, Corbacioglu S. 2009. Clinical implications of *c-Kit* mutations in acute myelogenous leukemia. *Curr Hematol Malig Rep* **4**: 77–82. doi:10.1007/s11899-009-0011-8
- Marcucci G, Metzler KH, Schwind S, Becker H, Maharry K, Mrózek K, Radmacher MD, Kohlschmidt J, Nicolet D, Whitman SP, et al. 2012. Age-related prognostic impact of different types of *DNMT3A* mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J Clin Oncol* **30**: 742–750. doi:10.1200/JCO.2011.39.2092
- Mardis ER, Ding L, Dooling DJ, Larson DE, McLellan MD, Chen K, Koboldt DC, Fulton RS, Delehaunty KD, McGrath SD, et al. 2009. Recurring mutations found by sequencing an acute myeloid leukemia genome. *New Engl J Med* **361**: 1058–1066. doi:10.1056/NEJMoa0903840
- Maxson JE, Ries RE, Wang YC, Gerbing RB, Kolb EA, Thompson SL, Guidry Auvil JM, Marra MA, Ma Y, Zong Z, et al. 2016. *CSF3R* mutations have a high degree of overlap with *CEBPA* mutations in pediatric AML. *Blood* **127**: 3094–3098. doi:10.1182/blood-2016-04-709899
- Mazumdar C, Shen Y, Xavy S, Zhao F, Reinisch A, Li R, Corces MR, Flynn RA, Buenrostro JD, Chan SM, et al. 2015. Leukemia-associated cohesin mutants dominantly enforce stem cell programs and impair human hematopoietic progenitor differentiation. *Cell Stem Cell* **17**: 675–688. doi:10.1016/j.stem.2015.09.017
- McMahon CM, Ferng T, Canaani J, Wang ES, Morrisette JJD, Eastburn DJ, Pellegrino M, Durruthy-Durruthy R, Watt CD, Asthana S, et al. 2019. Clonal selection with RAS pathway activation mediates secondary clinical resistance to selective FLT3 inhibition in acute myeloid leukemia. *Cancer Discov* **9**: 1050–1063. doi:10.1158/2159-8290.CD-18-1453
- McNerney ME, Brown CD, Wang X, Bartom ET, Karmakar S, Bandlamudi C, Yu S, Ko J, Sandall BP, Stricker T, et al. 2013. *CUX1* is a haploinsufficient tumor suppressor gene on chromosome 7 frequently inactivated in acute myeloid leukemia. *Blood* **121**: 975–983. doi:10.1182/blood-2012-04-426965
- Medeiros BC, Fathi AT, DiNardo CD, Pollyea DA, Chan SM, Swords R. 2017. Isocitrate dehydrogenase mutations in myeloid malignancies. *Leukemia* **31**: 272–281. doi:10.1038/leu.2016.275

A. Kishtagari and R.L. Levine

- Meisel M, Hinterleitner R, Pacis A, Chen L, Earley ZM, Mayassi T, Pierre JF, Ernest JD, Galipeau HJ, Thuille N, et al. 2018. Microbial signals drive pre-leukaemic myeloproliferation in a *Tet2*-deficient host. *Nature* **557**: 580–584. doi:10.1038/s41586-018-0125-z
- Mendler JH, Maharry K, Radmacher MD, Mrózek K, Becker H, Metzeler KH, Schwind S, Whitman SP, Khalife J, Kohlschmidt J, et al. 2012. *RUNX1* mutations are associated with poor outcome in younger and older patients with cytogenetically normal acute myeloid leukemia and with distinct gene and MicroRNA expression signatures. *J Clin Oncol* **30**: 3109–3118. doi:10.1200/JCO.2011.40.6652
- Meshinchi S, Appelbaum FR. 2009. Structural and functional alterations of *FLT3* in acute myeloid leukemia. *Clin Cancer Res* **15**: 4263–4269. doi:10.1158/1078-0432.CCR-08-1123
- Metzeler KH, Becker H, Maharry K, Radmacher MD, Kohlschmidt J, Mrózek K, Nicolet D, Whitman SP, Wu YZ, Schwind S, et al. 2011. *ASXL1* mutations identify a high-risk subgroup of older patients with primary cytogenetically normal AML within the ELN Favorable genetic category. *Blood* **118**: 6920–6929. doi:10.1182/blood-2011-08-368225
- Meyer C, Burmeister T, Gröger D, Tsaur G, Fechina L, Renneville A, Sutton R, Venn NC, Emerenciano M, Pombo-de-Oliveira MS, et al. 2018. The *MLL* recombinome of acute leukemias in 2017. *Leukemia* **32**: 273–284. doi:10.1038/leu.2017.213
- Micol JB, Abdel-Wahab O. 2016. The role of additional sex combs-like proteins in cancer. *Cold Spring Harb Perspect Med* **6**: a026526. doi:10.1101/cshperspect.a026526
- Mohapatra B, Ahmad G, Nadeau S, Zutshi N, An W, Scheffe S, Dong L, Feng D, Goetz B, Arya P, et al. 2013. Protein tyrosine kinase regulation by ubiquitination: Critical roles of Cbl-family ubiquitin ligases. *Biochim Biophys Acta* **1833**: 122–139. doi:10.1016/j.bbamer.2012.10.010
- Mullenders J, Aranda-Orgilles B, Lhoumaud P, Keller M, Pae J, Wang K, Kayembe C, Rocha PP, Raviram R, Gong Y, et al. 2015. Cohesin loss alters adult hematopoietic stem cell homeostasis, leading to myeloproliferative neoplasms. *J Exp Med* **212**: 1833–1850. doi:10.1084/jem.20151323
- Nagamachi A, Matsui H, Asou H, Ozaki Y, Aki D, Kanai A, Takubo K, Suda T, Nakamura T, Wolff L, et al. 2013. Haploinsufficiency of *SAMD9L*, an endosome fusion facilitator, causes myeloid malignancies in mice mimicking human diseases with monosomy 7. *Cancer Cell* **24**: 305–317. doi:10.1016/j.ccr.2013.08.011
- Nasmyth K, Haering CH. 2009. Cohesin: Its roles and mechanisms. *Annu Rev Genet* **43**: 525–558. doi:10.1146/annurev-genet-102108-134233
- Obeng EA, Chappell RJ, Seiler M, Chen MC, Campagna DR, Schmidt PJ, Schneider RK, Lord AM, Wang L, Gambe RG, et al. 2016. Physiologic expression of *Sf3b1*^{K700E} causes impaired erythropoiesis, aberrant splicing, and sensitivity to therapeutic spliceosome modulation. *Cancer Cell* **30**: 404–417. doi:10.1016/j.ccell.2016.08.006
- Okano M, Xie S, Li E. 1998. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet* **19**: 219–220. doi:10.1038/890
- Ostergaard P, Simpson MA, Connell FC, Steward CG, Brice G, Woollard WJ, Dafou D, Kilo T, Smithson S, Lunt P, et al. 2011. Mutations in *GATA2* cause primary lymphedema associated with a predisposition to acute myeloid leukemia (Emberger syndrome). *Nat Genet* **43**: 929–931. doi:10.1038/ng.923
- Pabst T, Mueller BU, Zhang P, Radomska HS, Narravula S, Schnittger S, Behre G, Hiddemann W, Tenen DG. 2001. Dominant-negative mutations of *CEBPA*, encoding CCAAT/enhancer binding protein- α (C/EBP α), in acute myeloid leukemia. *Nat Genet* **27**: 263–270. doi:10.1038/85820
- Pan F, Wingo TS, Zhao Z, Gao R, Makishima H, Qu G, Lin L, Yu M, Ortega JR, Wang J, et al. 2017. *Tet2* loss leads to hypermutagenicity in haematopoietic stem/progenitor cells. *Nat Commun* **8**: 15102. doi:10.1038/ncomms15102
- Papaemmanuil E, Gerstung M, Bullinger L, Gaidzik VI, Paschka P, Roberts ND, Potter NE, Heuser M, Thol F, Bolli N, et al. 2016. Genomic classification and prognosis in acute myeloid leukemia. *New Engl J Med* **374**: 2209–2221. doi:10.1056/NEJMoa1516192
- Parkin B, Ouilllette P, Wang Y, Liu Y, Wright W, Roulston D, Purkayastha A, Dressel A, Karp J, Bockenstedt P, et al. 2010. NF1 inactivation in adult acute myelogenous leukemia. *Clinical Cancer Res* **16**: 4135–4147. doi:10.1158/1078-0432.CCR-09-2639
- Paschka P, Marcucci G, Ruppert AS, Mrózek K, Chen H, Kittles RA, Vukosavljevic T, Perrotti D, Vardiman JW, Carroll AJ, et al. 2006. Adverse prognostic significance of *KIT* mutations in adult acute myeloid leukemia with inv(16) and t(8;21): A cancer and Leukemia Group B Study. *J Clin Oncol* **24**: 3904–3911. doi:10.1200/JCO.2006.06.9500
- Patel JP, Gönen M, Figueroa ME, Fernandez H, Sun Z, Racevskis J, Van Vlierberghe P, Dolgalev I, Thomas S, Aminova O, et al. 2012. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *New Engl J Med* **366**: 1079–1089. doi:10.1056/NEJMoa1112304
- Perl AE, Martinelli G, Cortes JE, Neubauer A, Berman E, Paolini S, Montesinos P, Baer MR, Larson RA, Ustun C, et al. 2019. Abstract CT184: Gilteritinib significantly prolongs overall survival in patients with *FLT3*-mutated (*FLT3*^{mut}) relapsed/refractory (R/R) acute myeloid leukemia (AML): Results from the Phase III ADMIRAL trial. *Cancer Res* **79**: CT184.
- Pollyea DA, Stevens BM, Jones CL, Winters A, Pei S, Minhajuddin M, D'Alessandro A, Culp-Hill R, Riemondy KA, Gillen AE, et al. 2018. Venetoclax with azacitidine disrupts energy metabolism and targets leukemia stem cells in patients with acute myeloid leukemia. *Nat Med* **24**: 1859–1866. doi:10.1038/s41591-018-0233-1
- Pronier E, Bowman RL, Ahn J, Glass J, Kandath C, Merlinsky TR, Whitfield JT, Durham BH, Gruet A, Hanasoge Somasundara AV, et al. 2018. Genetic and epigenetic evolution as a contributor to WT1-mutant leukemogenesis. *Blood* **132**: 1265–1278. doi:10.1182/blood-2018-03-837468
- Quek L, David MD, Kennedy A, Metzner M, Amatangelo M, Shih A, Stoilova B, Quivoron C, Heiblig M, Willekens C, et al. 2018. Clonal heterogeneity of acute myeloid leukemia treated with the IDH2 inhibitor enasidenib. *Nat Med* **24**: 1167–1177. doi:10.1038/s41591-018-0115-6
- Raffel S, Falcone M, Kneisel N, Hansson J, Wang W, Lutz C, Bullinger L, Poschet G, Nonnenmacher Y, Barnert A, et al.

2017. BCAT1 restricts α KG levels in AML stem cells leading to IDH^{mut}-like DNA hypermethylation. *Nature* **551**: 384–388. doi:10.1038/nature24294
- Rampal R, Figueroa ME. 2016. Wilms tumor 1 mutations in the pathogenesis of acute myeloid leukemia. *Haematologica* **101**: 672–679. doi:10.3324/haematol.2015.141796
- Rampal R, Alkalin A, Madzo J, Vasanthakumar A, Pronier E, Patel J, Li Y, Ahn J, Abdel-Wahab O, Shih A, et al. 2014. DNA hydroxymethylation profiling reveals that *WT1* mutations result in loss of TET2 function in acute myeloid leukemia. *Cell Rep* **9**: 1841–1855. doi:10.1016/j.celrep.2014.11.004
- Rogers HJ, Vardiman JW, Anastasi J, Raca G, Savage NM, Cherry AM, Arber D, Moore E, Morrisette JJ, Bagg A, et al. 2014. Complex or monosomal karyotype and not blast percentage is associated with poor survival in acute myeloid leukemia and myelodysplastic syndrome patients with inv(3)(q21q26.2)/t(3;3)(q21;q26.2): A Bone Marrow Pathology Group study. *Haematologica* **99**: 821–829. doi:10.3324/haematol.2013.096420
- Russler-Germain DA, Spencer DH, Young MA, Lamprecht TL, Miller CA, Fulton R, Meyer MR, Erdmann-Gilmore P, Townsend RR, Wilson RK, et al. 2014. The R882H DNMT3A mutation associated with AML dominantly inhibits wild-type DNMT3A by blocking its ability to form active tetramers. *Cancer Cell* **25**: 442–454. doi:10.1016/j.ccr.2014.02.010
- Saito S, Cigdem S, Okuwaki M, Nagata K. 2016. Leukemia-associated Nup214 fusion proteins disturb the XPO1-mediated nuclear-cytoplasmic transport pathway and thereby the NF- κ B signaling pathway. *Mol Cell Biol* **36**: 1820–1835. doi:10.1128/MCB.00158-16
- Sanada M, Suzuki T, Shih LY, Otsu M, Kato M, Yamazaki S, Tamura A, Honda H, Sakata-Yanagimoto M, Kumano K, et al. 2009. Gain-of-function of mutated *C-BL* tumour suppressor in myeloid neoplasms. *Nature* **460**: 904–908. doi:10.1038/nature08240
- Scheuermann JC, de Ayala Alonso AG, Oktaba K, Ly-Hartig N, McGinty RK, Fraterman S, Wilms M, Muir TW, Müller J. 2010. Histone H2A deubiquitinase activity of the Polycomb repressive complex PR-DUB. *Nature* **465**: 243–247. doi:10.1038/nature08966
- Schnittger S, Eder C, Jeromin S, Alpermann T, Fasan A, Grossmann V, Kohlmann A, Illig T, Klopp N, Wichmann HE, et al. 2013. *ASXL1* exon 12 mutations are frequent in AML with intermediate risk karyotype and are independently associated with an adverse outcome. *Leukemia* **27**: 82–91. doi:10.1038/leu.2012.262
- Schubbert S, Shannon K, Bollag G. 2007. Hyperactive Ras in developmental disorders and cancer. *Nat Rev Cancer* **7**: 295–308. doi:10.1038/nrc2109
- Schuurhuis GJ, Heuser M, Freeman S, Béné MC, Buccisano F, Cloos J, Grimwade D, Haferlach T, Hills RK, Hourigan CS, et al. 2018. Minimal/measurable residual disease in AML: A consensus document from the European LeukemiaNet MRD Working Party. *Blood* **131**: 1275–1291. doi:10.1182/blood-2017-09-801498
- Seiler M, Yoshimi A, Darman R, Chan B, Keaney G, Thomas M, Agrawal AA, Caleb B, Csibi A, Sean E, et al. 2018. H3B-8800, an orally available small-molecule splicing modulator, induces lethality in spliceosome-mutant cancers. *Nat Med* **24**: 497–504. doi:10.1038/nm.4493
- Sengsayadeth SM, Jagasia M, Engelhardt BG, Kassim A, Strickland SA, Goodman S, Lucid C, Vnencak-Jones CL, Greer JP, Savani BN. 2012. Allo-SCT for high-risk AML-CR1 in the molecular era: Impact of FLT3/ITD outweighs the conventional markers. *Bone Marrow Transplant* **47**: 1535–1537. doi:10.1038/bmt.2012.88
- Shih AH, Jiang Y, Meydan C, Shank K, Pandey S, Barreiro L, Antony-Debre I, Viale A, Socci N, Sun Y, et al. 2015. Mutational cooperativity linked to combinatorial epigenetic gain of function in acute myeloid leukemia. *Cancer Cell* **27**: 502–515. doi:10.1016/j.ccell.2015.03.009
- Shlush LI, Zandi S, Mitchell A, Chen WC, Brandwein JM, Gupta V, Kennedy JA, Schimmer AD, Schuh AC, Yee KW, et al. 2014. Identification of pre-leukaemic haematopoietic stem cells in acute leukaemia. *Nature* **506**: 328–333. doi:10.1038/nature13038
- Shlush LI, Mitchell A, Heisler L, Abelson S, Ng SWK, Trotman-Grant A, Medeiros JF, Rao-Bhatia A, Jaciw-Zurawski I, Marke R, et al. 2017. Tracing the origins of relapse in acute myeloid leukaemia to stem cells. *Nature* **547**: 104–108. doi:10.1038/nature22993
- Siegel RL, Miller KD, Jemal A. 2020. Cancer statistics, 2020. *CA Cancer J Clin* **70**: 7–30. doi:10.3322/caac.21590
- Sloand EM, Pfannes L, Chen G, Shah S, Solomou EE, Barrett J, Young NS. 2007. CD34 cells from patients with trisomy 8 myelodysplastic syndrome (MDS) express early apoptotic markers but avoid programmed cell death by up-regulation of antiapoptotic proteins. *Blood* **109**: 2399–2405. doi:10.1182/blood-2006-01-030643
- Smith ML, Cavenagh JD, Lister TA, Fitzgibbon J. 2004. Mutation of *CEBPA* in familial acute myeloid leukemia. *New Engl J Med* **351**: 2403–2407. doi:10.1056/NEJMoa041331
- Song WJ, Sullivan MG, Legare RD, Hutchings S, Tan X, Kufrin D, Ratajczak J, Resende IC, Haworth C, Hock R, et al. 1999. Haploinsufficiency of *CBFA2* causes familial thrombocytopenia with propensity to develop acute myelogenous leukaemia. *Nat Genet* **23**: 166–175. doi:10.1038/13793
- Sportoletti P, Varasano E, Rossi R, Mupo A, Tiacchi E, Vassiliou G, Martelli MP, Falini B. 2015. Mouse models of *NPM1*-mutated acute myeloid leukemia: Biological and clinical implications. *Leukemia* **29**: 269–278. doi:10.1038/leu.2014.257
- Stein EM, DiNardo CD, Pollyea DA, Fathi AT, Roboz GJ, Altman JK, Stone RM, DeAngelo DJ, Levine RL, Flinn IW, et al. 2017. Enasidenib in mutant *IDH2* relapsed or refractory acute myeloid leukemia. *Blood* **130**: 722–731. doi:10.1182/blood-2017-04-779405
- Stein EM, Garcia-Manero G, Rizzieri DA, Tibes R, Berdeja JG, Savona MR, Jongen-Lavrenic M, Altman JK, Thomson B, Blakemore SJ, et al. 2018. The DOT1L inhibitor pinometostat reduces H3K79 methylation and has modest clinical activity in adult acute leukemia. *Blood* **131**: 2661–2669. doi:10.1182/blood-2017-12-818948
- Stengel A, Kern W, Haferlach T, Meggendorfer M, Fasan A, Haferlach C. 2017. The impact of *TP53* mutations and *TP53* deletions on survival varies between AML, ALL, MDS and CLL: An analysis of 3307 cases. *Leukemia* **31**: 705–711. doi:10.1038/leu.2016.263
- Steudel C, Wermke M, Schaich M, Schäkel U, Illmer T, Ehninger G, Thiede C. 2003. Comparative analysis of *MLL* partial tandem duplication and *FLT3* internal tandem

A. Kishtagari and R.L. Levine

- duplication mutations in 956 adult patients with acute myeloid leukemia. *Genes Chromosomes Cancer* **37**: 237–251. doi:10.1002/gcc.10219
- Stone RM, Mandrekar SJ, Sanford BL, Laumann K, Geyer S, Bloomfield CD, Thiede C, Prior TW, Döhner K, Marcucci G, et al. 2017. Midostaurin plus chemotherapy for acute myeloid leukemia with a *FLT3* mutation. *New Engl J Med* **377**: 454–464. doi:10.1056/NEJMoa1614359
- Strissel PL, Strick R, Rowley JD, Zeleznik-Le NJ. 1998. An in vivo topoisomerase II cleavage site and a DNase I hypersensitive site colocalize near exon 9 in the MLL breakpoint cluster region. *Blood* **92**: 3793–3803. doi:10.1182/blood.V92.10.3793
- Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, Brudno Y, Agarwal S, Iyer LM, Liu DR, Aravind L, et al. 2009. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**: 930–935. doi:10.1126/science.1170116
- Tahirou TH, Inoue-Bungo T, Morii H, Fujikawa A, Sasaki M, Kimura K, Shiina M, Sato K, Kumasaka T, Yamamoto M, et al. 2001. Structural analyses of DNA recognition by the AML1/Runx-1 Runt domain and its allosteric control by CBF β . *Cell* **104**: 755–767. doi:10.1016/S0092-8674(01)00271-9
- Tartaglia M, Niemeyer CM, Fragale A, Song X, Buechner J, Jung A, Hählen K, Hasle H, Licht JD, Gelb BD. 2003. Somatic mutations in *PTPN11* in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia. *Nat Genet* **34**: 148–150. doi:10.1038/ng1156
- Taylor SJ, Thien CB, Dagger SA, Duyvestyn JM, Grove CS, Lee BH, Gilliland DG, Langdon WY. 2015. Loss of c-Cbl E3 ubiquitin ligase activity enhances the development of myeloid leukemia in FLT3-ITD mutant mice. *Exp Hematol* **43**: 191–206.e1. doi:10.1016/j.exphem.2014.11.009
- Thiede C, Steudel C, Mohr B, Schaich M, Schäkel U, Platzbecker U, Wermke M, Bornhäuser M, Ritter M, Neubauer A, et al. 2002. Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: Association with FAB subtypes and identification of subgroups with poor prognosis. *Blood* **99**: 4326–4335. doi:10.1182/blood.V99.12.4326
- Thota S, Viny AD, Makishima H, Spitzer B, Radivoyevitch T, Przychodzen B, Sekeres MA, Levine RL, Maciejewski JP. 2014. Genetic alterations of the cohesin complex genes in myeloid malignancies. *Blood* **124**: 1790–1798. doi:10.1182/blood-2014-04-567057
- Todd MA, Picketts DJ. 2012. PHF6 interacts with the nucleosome remodeling and deacetylation (NuRD) complex. *J Proteome Res* **11**: 4326–4337. doi:10.1021/pr3004369
- Tsai FY, Keller G, Kuo FC, Weiss M, Chen J, Rosenblatt M, Alt FW, Orkin SH. 1994. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature* **371**: 221–226. doi:10.1038/371221a0
- Tyner JW, Tognon CE, Bottomly D, Wilmot B, Kurtz SE, Savage SL, Long N, Schultz AR, Traer E, Abel M, et al. 2018. Functional genomic landscape of acute myeloid leukaemia. *Nature* **562**: 526–531. doi:10.1038/s41586-018-0623-z
- Uckelmann HJ, Kim SM, Wong EM, Hatton C, Giovinozzo H, Gadrey JY, Krivtsov AV, Rücker FG, Döhner K, McGeehan GM, et al. 2020. Therapeutic targeting of pre-leukemia cells in a mouse model of *NPM1* mutant acute myeloid leukemia. *Science* **367**: 586–590. doi:10.1126/science.aax5863
- Van Vlierberghe P, Patel J, Abdel-Wahab O, Lobry C, Hedvat CV, Balbin M, Nicolas C, Payer AR, Fernandez HF, Tallman MS, et al. 2011. *PHF6* mutations in adult acute myeloid leukemia. *Leukemia* **25**: 130–134. doi:10.1038/leu.2010.247
- Viny AD, Ott CJ, Spitzer B, Rivas M, Meydan C, Papalexi E, Yelin D, Shank K, Reyes J, Chiu A, et al. 2015. Dose-dependent role of the cohesin complex in normal and malignant hematopoiesis. *J Exp Med* **212**: 1819–1832. doi:10.1084/jem.20151317
- Viny AD, Bowman RL, Liu Y, Lavallée VP, Eisman SE, Xiao W, Durham BH, Navitski A, Park J, Braunstein S, et al. 2019. Cohesin members Stag1 and Stag2 display distinct roles in chromatin accessibility and topological control of HSC self-renewal and differentiation. *Cell Stem Cell* **25**: 682–696.e8. doi:10.1016/j.stem.2019.08.003
- Wang J, Fernald AA, Anastasi J, Le Beau MM, Qian Z. 2010. Haploinsufficiency of *Apc* leads to ineffective hematopoiesis. *Blood* **115**: 3481–3488. doi:10.1182/blood-2009-11-251835
- Wang Y, Xiao M, Chen X, Chen L, Xu Y, Lv L, Wang P, Yang H, Ma S, Lin H, et al. 2015. WT1 recruits TET2 to regulate its target gene expression and suppress leukemia cell proliferation. *Mol Cell* **57**: 662–673. doi:10.1016/j.molcel.2014.12.023
- Wang E, Lu SX, Pastore A, Chen X, Imig J, Chun-Wei Lee S, Hockemeyer K, Ghebrechristos YE, Yoshimi A, Inoue D, et al. 2019. Targeting an RNA-binding protein network in acute myeloid leukemia. *Cancer Cell* **35**: 369–384.e7. doi:10.1016/j.ccell.2019.01.010
- Welch JS, Ley TJ, Link DC, Miller CA, Larson DE, Koboldt DC, Wartman LD, Lamprecht TL, Liu F, Xia J, et al. 2012. The origin and evolution of mutations in acute myeloid leukemia. *Cell* **150**: 264–278. doi:10.1016/j.cell.2012.06.023
- Wendorff AA, Quinn SA, Rashkovan M, Madubata CJ, Ambesi-Impiombato A, Litzow MR, Tallman MS, Paietta E, Paganin M, Basso G, et al. 2019. *Phf6* loss enhances HSC self-renewal driving tumor initiation and leukemia stem cell activity in T-ALL. *Cancer Discov* **9**: 436–451. doi:10.1158/2159-8290.CD-18-1005
- Wendt KS, Yoshida K, Itoh T, Bando M, Koch B, Schirghuber E, Tsutsumi S, Nagae G, Ishihara K, Mishihiro T, et al. 2008. Cohesin mediates transcriptional insulation by CCCTC-binding factor. *Nature* **451**: 796–801. doi:10.1038/nature06634
- West RR, Hsu AP, Holland SM, Cuellar-Rodriguez J, Hickstein DD. 2014. Acquired *ASXL1* mutations are common in patients with inherited *GATA2* mutations and correlate with myeloid transformation. *Haematologica* **99**: 276–281. doi:10.3324/haematol.2013.090217
- Wong TN, Ramsingh G, Young AL, Miller CA, Touma W, Welch JS, Lamprecht TL, Shen D, Hundal J, Fulton RS, et al. 2015. Role of *TP53* mutations in the origin and evolution of therapy-related acute myeloid leukaemia. *Nature* **518**: 552–555. doi:10.1038/nature13968
- Wouters BJ, Löwenberg B, Erpelinck-Verschueren CA, van Putten WL, Valk PJ, Delwel R. 2009. Double *CEBPA* mu-

Somatic Mutations in Acute Myeloid Leukemia

- tations, but not single *CEBPA* mutations, define a subgroup of acute myeloid leukemia with a distinctive gene expression profile that is uniquely associated with a favorable outcome. *Blood* **113**: 3088–3091. doi:10.1182/blood-2008-09-179895
- Xie M, Lu C, Wang J, McLellan MD, Johnson KJ, Wendl MC, McMichael JF, Schmidt HK, Yellapantula V, Miller CA, et al. 2014. Age-related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med* **20**: 1472–1478. doi:10.1038/nm.3733
- Xu W, Yang H, Liu Y, Yang Y, Wang P, Kim SH, Ito S, Yang C, Wang P, Xiao MT, et al. 2011. Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of α -ketoglutarate-dependent dioxygenases. *Cancer Cell* **19**: 17–30. doi:10.1016/j.ccr.2010.12.014
- Yarden Y, Kuang WJ, Yang-Feng T, Coussens L, Munemitsu S, Dull TJ, Chen E, Schlessinger J, Francke U, Ullrich A. 1987. Human proto-oncogene *c-kit*: A new cell surface receptor tyrosine kinase for an unidentified ligand. *EMBO J* **6**: 3341–3351. doi:10.1002/j.1460-2075.1987.tb02655.x
- Yokoyama A, Somervaille TC, Smith KS, Rozenblatt-Rosen O, Meyerson M, Cleary ML. 2005. The menin tumor suppressor protein is an essential oncogenic cofactor for MLL-associated leukemogenesis. *Cell* **123**: 207–218. doi:10.1016/j.cell.2005.09.025
- Yoshida K, Sanada M, Shiraishi Y, Nowak D, Nagata Y, Yamamoto R, Sato Y, Sato-Otsubo A, Kon A, Nagasaki M, et al. 2011. Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature* **478**: 64–69. doi:10.1038/nature10496
- Yoshimi A, Lin KT, Wiseman DH, Rahman MA, Pastore A, Wang B, Lee SC, Micol JB, Zhang XJ, de Botton S, et al. 2019. Coordinated alterations in RNA splicing and epigenetic regulation drive leukaemogenesis. *Nature* **574**: 273–277. doi:10.1038/s41586-019-1618-0
- Yoshizato T, Dumitriu B, Hosokawa K, Makishima H, Yoshida K, Townsley D, Sato-Otsubo A, Sato Y, Liu D, Suzuki H, et al. 2015. Somatic mutations and clonal hematopoiesis in aplastic anemia. *New Engl J Med* **373**: 35–47. doi:10.1056/NEJMoa1414799
- Zhang Q, Zhao K, Shen Q, Han Y, Gu Y, Li X, Zhao D, Liu Y, Wang C, Zhang X, et al. 2015. Tet2 is required to resolve inflammation by recruiting Hdac2 to specifically repress IL-6. *Nature* **525**: 389–393. doi:10.1038/nature15252
- Zhou L, Opalinska J, Sohal D, Yu Y, Mo Y, Bhagat T, Abdel-Wahab O, Fazzari M, Figueroa M, Alencar C, et al. 2011. Aberrant epigenetic and genetic marks are seen in myelodysplastic leukocytes and reveal *Dock4* as a candidate pathogenic gene on chromosome 7q. *J Biol Chem* **286**: 25211–25223. doi:10.1074/jbc.M111.235028

The Role of Somatic Mutations in Acute Myeloid Leukemia Pathogenesis

Ashwin Kishtagari and Ross L. Levine

Cold Spring Harb Perspect Med 2021; doi: 10.1101/cshperspect.a034975 originally published online May 12, 2020

Subject Collection [Leukemia and Lymphoma: Molecular and Therapeutic Insights](#)

Oncogenic Mechanisms and Therapeutic Targeting of Metabolism in Leukemia and Lymphoma

Maximilian Stahl, Zachary D. Epstein-Peterson and Andrew M. Intlekofer

Non-Hodgkin Lymphomas: Malignancies Arising from Mature B Cells

Jennifer Shingleton, Jie Wang, Carolyn Baloh, et al.

MAP-Kinase-Driven Hematopoietic Neoplasms: A Decade of Progress in the Molecular Age

Rikhia Chakraborty, Omar Abdel-Wahab and Benjamin H. Durham

Mouse Models in the Study of Mature B-Cell Malignancies

Laura Pasqualucci and Ulf Klein

Mouse Models of Myeloid Malignancies

Faisal Basheer and George Vassiliou

Epigenetic Mechanisms in Leukemias and Lymphomas

Cihangir Duy, Wendy Béguelin and Ari Melnick

The Biology of B-Progenitor Acute Lymphoblastic Leukemia

Kathryn G. Roberts and Charles G. Mullighan

RNA Regulators in Leukemia and Lymphoma

Camila Prieto and Michael G. Kharas

Biological and Translational Considerations regarding the Recent Therapeutic Successes and Upcoming Challenges for Multiple Myeloma

Constantine S. Mitsiades

Biology and Molecular Pathogenesis of Mature T-Cell Lymphomas

José R. Cortés and Teresa Palomero

The Role of Somatic Mutations in Acute Myeloid Leukemia Pathogenesis

Ashwin Kishtagari and Ross L. Levine

Chronic Lymphocytic Leukemia

Nicholas Chiorazzi, Shih-Shih Chen and Kanti R. Rai

Normal Hematopoiesis Is a Balancing Act of Self-Renewal and Regeneration

Oakley C. Olson, Yoon-A Kang and Emmanuelle Passegué

Impact of Genetics on Mature Lymphoid Leukemias and Lymphomas

Nathanael G. Bailey and Kojo S.J. Elenitoba-Johnson

Epidemiology and Etiology of Leukemia and Lymphoma

Jordan A. Baeker Bispo, Paulo S. Pinheiro and Erin K. Kobetz

Clonal Hematopoiesis and Premalignant Diseases

Justin Kaner, Pinkal Desai, Nuria Mencia-Trinchant, et al.

For additional articles in this collection, see <http://perspectivesinmedicine.cshlp.org/cgi/collection/>