The Spectrum of Fungi That Infects Humans

Julia R. Köhler¹, Arturo Casadevall², and John Perfect³

¹Division of Infectious Diseases, Children's Hospital, Harvard Medical School, Boston, Massachusetts 02115

²Departments of Microbiology and Immunology and Medicine, Division of Infectious Diseases, Albert Einstein College of Medicine, New York, New York 10461

³Division of Infectious Diseases, Duke Medical Center, Durham, North Carolina 27710

Correspondence: julia.koehler@childrens.harvard.edu

Few among the millions of fungal species fulfill four basic conditions necessary to infect humans: high temperature tolerance, ability to invade the human host, lysis and absorption of human tissue, and resistance to the human immune system. In previously healthy individuals, invasive fungal disease is rare because animals' sophisticated immune systems evolved in constant response to fungal challenges. In contrast, fungal diseases occur frequently in immunocompromised patients. Paradoxically, successes of modern medicine have put increasing numbers of patients at risk for invasive fungal infections. Uncontrolled HIV infection additionally makes millions vulnerable to lethal fungal diseases. A concerted scientific and social effort is needed to meet these challenges.

Fungal infections today are among the most difficult diseases to manage in humans. Some fungi cause disease in healthy people, but most fungal infections occur in individuals already experiencing serious illness, and frequently jeopardize the success of the newest medical advances in cancer care, solid organ and hematopoietic stem cell transplantation, neonatal medicine, autoimmune disease therapies, trauma and intensive care, and sophisticated surgery. In fact, these medical advances themselves often make their beneficiaries vulnerable to fungal disease. Treatment can involve breaching normal immune functions, or susceptible patients, such as extremely premature newborns who survive long enough to become infected by a fungus.

The following discussion intends to touch on highlights of the evolutionary developments

by which living humans became substrates for fungi. Given the tremendous wealth of recent findings on fungal evolution, phylogenetics, genomics, development, and pathogenesis, this overview will necessarily omit much work critical to our understanding of fungi, which the other articles in this collection will focus on in detail.

Among the estimated 1.5–5.0 million fungal species on planet Earth (O'Brien et al. 2005), only several hundred cause disease in humans, and very few are able to affect healthy people. Animals coevolved with fungi, and the sophisticated and potent human immune system arose from the constant challenge posed by the microbial world. Fungal pathogens likely adapted their pathogenic repertoire to other animal prey—mammals, insects, and even unicellular amoebae—before encountering and attacking

Editors: Arturo Casadevall, Aaron P. Mitchell, Judith Berman, Kyung J. Kwon-Chung, John R. Perfect, and Joseph Heitman
Additional Perspectives on Human Fungal Pathogens available at www.perspectivesinmedicine.org

Copyright © 2015 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a019273
Cite this article as *Cold Spring Harb Perspect Med* 2015;5:a019273

J.R. Köhler et al.

humans. But unlike plants, insects, and ectothermic vertebrates (animals whose body temperature fluctuates with their surroundings), mammals are highly resistant to invasive fungal diseases, and evolution of endothermy and homeothermy enhanced antifungal immunity (Robert and Casadevall 2009; Bergman and Casadevall 2010). The remarkable resistance of mammals to fungal pathogens has been hypothesized to be responsible for emergence of mammals as the dominant land species, when proliferation of fungi at the end of the Cretaceous era created a “fungal filter” that selected for this animal group (Casadevall 2005, 2012).

For a fungus, parasitizing a human is a demanding strategy. Four criteria must be fulfilled. (1) It must be able to grow at a high temperature, at or above 37°C. (2) It must be able to reach the tissues it will parasitize, by penetrating host tissue barriers, or by circumventing them through small airborne cells that enter air-filled spaces of lungs and sinuses directly. (3) It must be able to digest and absorb components of human tissues. (4) Finally, it must be able to withstand the human immune system.

Virulence factors can be divided into aspects of physiology that allow a fungus to fulfill these four criteria. Growth at high temperature is a stringent criterion, because land-colonizing fungi likely evolved in association with plants (Humphreys et al. 2010), whose nighttime temperature must be cool enough to minimize the ratio of carbon-expending respiration to carbon-assimilating photosynthesis. Casadevall and colleagues have discussed how the stable elevated temperature of endothermic (warm-blooded) animals may have been one of the most potent developments in antifungal immunity (Robert and Casadevall 2009; Bergman and Casadevall 2010; Garcia-Solache and Casadevall 2010). The energy cost of the human fever response to infection, which is tightly regulated by cytokines like TNF- α , represents another testimony to the evolutionary importance of temperature as a major host defense against fungi.

The role of fungal heat intolerance is illustrated by the different clinical scenarios of disseminated *Fusarium* infection versus disseminated candidiasis. *Fusarium* spp. are important

plant pathogens that can infect neutropenic patients. As filamentous *Fusarium* spp. produce cells resembling yeast in the human host, which readily float within the bloodstream, a hallmark of their dissemination is development of numerous metastatic foci in the skin, presumably because the skin is the coolest organ. The development of metastatic lesions in warm organs distant from the primary focus, like spleen and kidneys, is much more unusual in fusariosis. In contrast, invasive infections with human temperature-adapted *Candida* spp., which also use yeast to disseminate through the bloodstream, frequently lead to innumerable metastatic foci in deep organs.

Morphogenesis is an important virulence factor related to fungal locomotion. Although fungal hyphae can pierce tissue barriers owing to the turgor pressure at their tips, yeast can readily disseminate to distant sites in a large animal. Fungi that infect healthy humans do so almost exclusively in their yeast form. But for most fungi, the ability to assume various shapes is critical for infecting humans, because many enter the body in the form of small, round airborne dispersal propagules, sporangiospores or conidia, which are produced from hyphal cells. In the oceans, fungi originated as unicellular, oval cells propelled by flagella, so growth as a round or oval cell was a fundamental fungal trait. Lucking et al. (2009), in their comparison of fossil and molecular dating systems of fungal evolution, state that “all dating estimates show that the evolution of filamentous fungi occurred much later than the origin of the fungal lineage itself, suggesting that for a long time after their origin, fungi were heterotrophic, unicellular, flagellate, aquatic organisms.” An option for the hyphal form, whose adaptive utility is shown by convergent evolution of hyphae in fungi and oomycete water molds (Money et al. 2004), may have evolved when fungi colonized land as plant symbionts (Redecker et al. 2000; Heckman et al. 2001; Lucking et al. 2009). Morphogenesis also contributes to protection against amoeboid cell predation. Filamentous forms of *Cryptococcus neoformans* are resistant to amoeba (Neilson et al. 1978), and incubation of *Histoplasma capsulatum* and *Blastomyces*

dermatitidis yeast with amoeba at 37°C triggers transition to filamentous growth (Steenbergen et al. 2004).

Tremendous variability of cell forms and the ability to switch between them continues in extant fungal phyla, and few lineages have completely lost a round or a filamentous cell form. A highly successful human-host-associated fungus, *Candida albicans*, is notable for the facility and frequency with which it switches between a broad spectrum of growth forms in the host, between round yeast, elongated pseudohyphae, and filamentous hyphae (Sudbery et al. 2004). Adhesion molecules are virulence factors related to fungal locomotion strategies because a hypha must anchor itself to host tissue to exert tissue-penetrating pressure at its tip, and because once a fungal cell has reached a favorable location, it needs to be able to stay at the site.

Secretion of digestive enzymes appropriate to dissolve host tissue is the first step for a fungus to use the host as a nutritious substrate. The next step is to absorb the small molecules released by digestion of host tissues; fungal cells have transporters of nitrogen and carbon sources as well as metal ions and other micronutrients. Iron acquisition plays a special role because of its importance in critical enzymatic processes including respiration, and because the human host actively sequesters iron during infection to withhold it from the pathogen (Boelaert et al. 1993; Ibrahim et al. 2006; Kronstad et al. 2013; Noble 2013). The importance of iron sequestration in the host to defend against fungal growth was dramatically illustrated by disseminated Mucorales infections in patients receiving the iron chelator, deferoxamine (Windus et al. 1987).

Most fungi, even with these capabilities, cannot withstand the phagocytes unleashed against them by an immunocompetent human, which have help from opsonins such as complement and specific antibody, and from activating T lymphocytes. Fungi that infect healthy humans devote a large portion of their physiology to withstanding or evading the immune system. To this end they use an amazingly diverse array of strategies, which sometimes differ even be-

The Spectrum of Fungi That Infects Humans

tween lineages of a single species. In contrast, many species that grow at 37°C can readily infect a severely immunocompromised patient.

Last, coordinating the processes to fulfill these four criteria of pathogenicity demands a complex network of sensing and signaling systems, which inform the fungal cell of external conditions and set in motion its appropriate responses. Loss of virulence was shown many times in many fungal pathogens, when just a single element of a sensing and signaling network was genetically or pharmacologically disturbed, such as mechanosensor, calcineurin, protein kinase A, MAP kinase, Tor, and high osmolarity signaling pathways (Rohde and Cardenas 2004; Cramer et al. 2008; Kumamoto 2008; Bastidas and Heitman 2009; Argimon et al. 2011; Ernst and Pla 2011; Hogan and Muhlischlegel 2011; Shapiro et al. 2012; Chen et al. 2013).

FUNGI THAT INFECT HEALTHY HUMANS

The capability to infect an immunocompetent human has arisen independently multiple times among three major fungal phyla: the Entomophthoromycota (Kwon-Chung 2012), the Ascomycota, and the Basidiomycota.

Entomophthoromycota

Entomophthoromycota contains effective pathogens of insects. They occur worldwide, but have been found as agents of invasive human infection only in subtropical and tropical regions. The species of genera pathogenic for humans, *Basidiobolus* and *Conidiobolus*, can be isolated from plant debris and soil especially during rainy months (Bittencourt 1988). As emphasized recently by Kwon-Chung (2012), these genera are evolutionarily distant from each other as well as from the Mucorales, with whom they traditionally were classified in Zygomycota.

Conidiobolus spp. cause submucosal disease of the nose, sinuses, and central face. In most cases, gradual progression causes swelling of submucosal and subcutaneous tissue, disfigurement, breathing difficulties, and chronic bacte-

J.R. Köhler et al.

rial sinusitis caused by the blockage of ostia. Unlike the Mucorales causing sinus disease, *Conidiobolus* does not usually invade blood vessels nor penetrate into the central nervous system (CNS) (Prabhu and Patel 2004).

Two species, *Conidiobolus coronatus* and *Conidiobolus incongruus*, have been isolated from central facial disease. Presumably, the route of infection is inhalation of large ballistospores forcibly launched from the sporangio-phores on which they are produced singly to distances up to 30 cm (Isa-Isa et al. 2012). Owing to their large size of 30–38 μm (Isa-Isa et al. 2012), spores presumably land on the mucosa of nasal air passages and fail to reach the distal airways (Bittencourt et al. 2006).

Conidiobolus spp. infect arthropods including mites, spiders, and insects (Isa-Isa et al. 2012). They elaborate lytic enzymes including elastases, collagenases, and lipases, likely to parasitize and kill arthropods (Comerio et al. 2008). These enzymes enable *Conidiobolus* to digest human tissue, growing by extension from nasal submucosa through the facial soft and bony tissue (Gugnani 1992). Metabolites toxic to insects have also been identified but because the fungus is highly lethal to arthropods and only of moderate virulence in mammals, it remains to be seen whether these metabolites play any role in human infection (Wieloch et al. 2011).

The host sends neutrophils, eosinophils, and histiocytes to contain the fungus, and granulomas (structured assemblies of histiocytes) develop. Histopathology can show dense eosinophilic material surrounding the hyphae, the Splendore-Hoeppli phenomenon, which has been suggested to consist of antigen–antibody precipitate (Isa-Isa et al. 2012).

Basidiobolus ranarum causes subcutaneous disease primarily in children of tropical and subtropical Africa, Asia, and the Americas. Nodules appear at sites of inoculation through insect bites, scratches, or small wounds (Ribes et al. 2000). In addition, *Basidiobolus* has caused gastrointestinal infections, presumably after ingestion of large inocula, and symptoms resemble inflammatory bowel disease (Zavasky et al. 1999; Vikram et al. 2012). When insects infected by *Basidiobolus* are eaten by amphibians and

reptiles, the fungus is shed in the feces (Ribes et al. 2000) and attaches to plant matter with which it is accidentally inoculated under the skin, for instance, by thorns, leaves used for cleansing, etc. As an insect pathogen, it also elaborates proteases and lipases (Echetebe and Ononogbu 1982; Okafor et al. 1987; Okafor and Gugnani 1990; Okafor 1994). Its inability to invade deep organs is likely related to limited thermotolerance; 37°C may be its maximal growth temperature (Ribes et al. 2000).

Ascomycota

Pathogenic Onygenales

Several soil-inhabiting members of the ascomycete order Onygenales have evolved to parasitize mammals and cause systemic infection. They have been classified in the family Ajellomycetaceae (Untereiner et al. 2004; Bagagli et al. 2008), which include *B. dermatitidis* (teleomorph, *Ajellomyces dermatitidis*), *H. capsulatum* (teleomorph, *Ajellomyces capsulatus*), *Paracoccidioides brasiliensis*, *Paracoccidioides lutzii*, *Lacazia loboi*, *Coccidioides immitis*, *Coccidioides posadasii*, and related fungi with similar properties. The intracellular pathogenic strategy of some of these organisms is similar in protozoa and macrophages (Steenbergen et al. 2001, 2003), raising the possibility that their capacity for virulence arose accidentally as a result of environmental pressures (Casadevall and Pirofski 2007).

Disease caused by these fungi begins asymptotically in the lungs and progresses to an influenza-like illness or frank pneumonia. They can disseminate to distant sites, can persist and reactivate, with different organ predilections among different genera: *Paracoccidioides* in oral and respiratory mucous membranes (Queiroz-Telles and Escuissato 2011; Marques 2012), *Blastomyces* in bones, joints, and skin (Bradsher et al. 2003), and *Histoplasma* in multiple organs including the gastrointestinal tract and adrenals, as well as bones and skin (Kauffman 2007).

Their primary lifestyle is saprobic. Most genera occur in defined geographic areas, which follow particular soil consistencies, for instance, dry and alkaline for the *Coccidioides* spp. and

acidic for *Histoplasma*. It has not been ruled out that their geographic distribution follows that of specific host mammals. In the soil, they grow as a branched network of hyphal filaments, a mycelium, and saprobically use organic matter. They produce airborne dispersal cells: Conidia produced from specialized hyphae, as *Histoplasma* and *Blastomyces* do, or arthrospores from regulated fragmentation of hyphal compartments, as *Coccidioides* and *Paracoccidioides* do. *Paracoccidioides* produces both arthroconidia and aleuriconidia (conidia directly produced along the hyphae). Once conidia or arthrospores are taken up by a mammal, usually by inhalation, they convert into parasitic yeast (or spherules, in the case of *Coccidioides* spp.) and initiate an infection. The principal signal for their conversion to the yeast form is the elevated mammalian body temperature, so that this switch can be induced in vitro when they are cultured at 37°C or above. For this reason they are grouped as thermal dimorphs. These fungi infect, persist, and cause disease in healthy hosts. In immunocompromised hosts, the infections are more clinically apparent, more severe, and more likely to disseminate to multiple organs.

Adaptive cellular immunity is required to control the infection with these fungi and can be detected by skin tests with purified antigens. For this reason, clinically apparent infections with some members of this group, such as *H. capsulatum*, became much more common with the rising prevalence of AIDS. These Onygenales (Ajellomycetaceae) are the only phylogenetically related group whose members all cause systemic disease in otherwise healthy people.

Other thermally dimorphic fungi are *Sporothrix schenckii* (order Ophiostomatales), which is primarily introduced into the host through skin injuries, and *Talaromyces* (*Penicillium*) *marneffei* (order Eurotiales), which does not infect healthy hosts but because of the AIDS epidemic has become a prevalent and important invasive pathogen in Southeast Asia. These fungi coevolved with higher vertebrates, as it is the adaptive immune system by which they are controlled. The fact that these evolutionarily distant fungi convert from saprobic hyphal to

pathogenic yeast growth at the temperatures of endothermic animals suggests that an ability to access large nitrogen and other nutrient stores of an animal host is an advantageous lifestyle option for a soil dweller, and a result of convergent evolution in several phyla.

H. capsulatum will be discussed at greater length because it is well studied and also because it exemplifies important principles for other thermally dimorphic Onygenales. This fungus inhabits soil and other organic matter such as decaying wood enriched with bat and bird droppings. Its hyphae produce macro- and microconidia, and the latter are small enough, <5 µm, to penetrate into the alveoli of mammalian lungs. There they are phagocytized by alveolar macrophages and converted to yeast that is able to proliferate in the hostile environment of the phagolysosome.

Aspects of histoplasmosis resemble infections with a well-adapted bacterial intracellular pathogen, *Mycobacterium tuberculosis*. Depending on inoculum size, infection is most often inapparent or leads to a self-limited influenza-like illness. The pathogen can persist indefinitely until immune decline allows it to reactivate. Lung disease resembling TB can ensue immediately after infection, and dissemination to all organs occurs depending on host immune status and age. The response of an immunocompetent host is granuloma formation.

Histoplasma occurs on all continents, although it is rare in Eurasia and most common in North and South America. Two studies of *H. capsulatum* showed that its growth and clinical phenotypes, which had traditionally been the basis for its division into varieties (*capsulatum*, *duboisii*, and *farcinosum*), did not align with the divisions of its seven phylogenetic clades (Kasuga et al. 1999, 2003), and that the different clinical disease states attributed to these (now obsolete) varieties were caused by the same clades.

Phylogenetic analyses suggest that *H. capsulatum* arose in South America 3.2–13.0 million years ago and its present distribution reflects a period of rapid spread to all continents except Eurasia (where it was later introduced), followed by restriction to the warmest areas during

J.R. Köhler et al.

a period of intense cold 1.8 million years ago (Kasuga et al. 1999, 2003). From there, it spread to temperate regions as the earth warmed again. This idea is consistent with its low genetic diversity in temperate regions and high diversity in tropical areas (Kasuga et al. 1999), mirroring the diversity of plant species in areas of temperate and tropical forest, which are its geographic areas of distribution.

The two North American clades are not closely related and are thought to have been genetically isolated for the past ~20 million years (Kasuga et al. 1999), despite overlapping territories. These two clades differ in virulence, with North American clade 1 having been isolated predominantly from AIDS patients, and clade 2 infecting immunocompetent hosts (Kasuga et al. 1999). A clade 2 isolate and a representative of the Panama clade have been used in genetic studies of virulence factors, and surprising differences were found, as discussed below.

The habitat of *Histoplasma* is soil enriched with bird and bat droppings, in specific geographic locations including the Ohio and Mississippi valleys of North America and moist regions of Central and South America. Whereas birds are very rarely infected with *H. capsulatum* (Quist et al. 2011), bats are frequently infected but respond with minimal inflammation (Taylor et al. 1999), suggesting active repression of the host immune response by the fungus. It was suggested that, “in at least some bat species natural exposure may result in a chronic, controlled infection, which allows the bat to excrete viable fungi over a long period” (Hoff and Bigler 1981). Experimentally infected bats also had a minimal inflammatory response (McMurray and Greer 1979).

A broad variety of other infected mammals such as cats (Aulakh et al. 2012), dogs (Brömel and Sykes 2005), sea otters (Morita et al. 2001), and badgers (Bauder et al. 2000) do respond with granulomatous inflammation to the presence of *Histoplasma* yeast. Perhaps *H. capsulatum* has evolved to infect bats, in which it avoids activation of host responses, by mechanisms only partially adapted to other immunocompetent mammals. If this is the case, the fungus has established an efficient amplification cycle by

which its mycelial form grows in droppings under bat roosting places, for instance, in caves, and its microconidia are released into the air to be inhaled by the bats that fertilize its soil substrate (Taylor et al. 1999).

Several mechanisms by which *H. capsulatum* yeast successfully interacts with mammalian hosts have been identified. Surprisingly, different clades use different strategies to achieve an intracellular lifestyle in macrophages, and to suppress macrophage activation. For example, macrophages recognize the important pathogen associated molecular pattern (PAMP) $\beta(1,3)$ -D-glucan with the receptor dectin-1 (Rappleye et al. 2007), and ligand binding triggers responses that enhance phagocyte activation responses (Underhill et al. 2005). $\beta(1,3)$ -D-glucan is the main structural polymer of ascomycete cell walls, so *Histoplasma* cannot forgo its use to avoid activating their hosts' macrophages. Instead, *Histoplasma* yeast of most clades hide it by coating their cell surface with $\alpha(1,3)$ -D-glucan. Perturbation of genes whose products are required for $\alpha(1,3)$ -D-glucan synthesis renders cells of the Panama clade avirulent (Rappleye et al. 2004). Notably, this $\alpha(1,3)$ -D-glucan coat is lacking in a North American clade 2 strain (Edwards et al. 2011), and an $\alpha(1,3)$ -D-glucan synthase mutant in this strain background is fully virulent (Edwards et al. 2011).

Another cell surface protein, Yps3, is expressed to a significant level only in the more virulent North American clade 2 (Bohse and Woods 2007). This protein, a homolog of *B. dermatitidis* BAD1, contributes to dissemination of *H. capsulatum* to extrapulmonary foci (Holbrook and Rappleye 2008). Its gene is present in most strains but substantial transcription is known to occur only in North American clade 2, the clade lacking $\alpha(1,3)$ -D-glucan. Yps3, like BAD1, may directly suppress TNF- α production, circumventing the need for this clade to use $\alpha(1,3)$ -D-glucan to hide the stimulatory PAMP $\beta(1,3)$ -D-glucan (Holbrook and Rappleye 2008). The known variations in their pathogenic repertoires among clades of *Histoplasma* are based in differences of regulatory rather than coding regions between the sequenced strains (Edwards et al. 2013).

H. capsulatum yeast have evolved a repertoire of manipulative molecules to turn the host compartment specifically evolved to kill pathogens, the phagolysosome, into a habitat in which they thrive (Sebghati et al. 2000; Youseff et al. 2009). Remarkably, within the teleomorph genus *Ajellomyces*, originally comprising the anamorph sister species *H. capsulatum* and *B. dermatitidis*, one species *H. capsulatum* chose this route of intracellular life, whereas the other, *B. dermatitidis*, is a successful extracellular pathogen.

Distinct *H. capsulatum* clades differ in specific details of the pathogenic program of their yeast, but they have in common that their suite of virulence factors is turned on simultaneously with the switch from hyphal to yeast growth. Mammalian body temperature of 37°C or above is the main signal to trigger conversion from the mycelium to yeast. This conversion entails differential transcription of ~20% of the genome (Nguyen and Sil 2008), and is controlled by a network of transcriptional regulators apparently responding to a histidine kinase signal (Nemecsek et al. 2006; Webster and Sil 2008; Cain et al. 2012; Beyhan et al. 2013). Mutants in these regulators are incapable of the transition to yeast growth, and are avirulent (Nemecsek et al. 2006; Webster and Sil 2008; Cain et al. 2012; Beyhan et al. 2013).

Other members of the *Ajellomycetaceae*, all of which associate with mammals, have equally distinctive ecology and pathobiology. Mycelia of the *Coccidioides* spp., growing near rodent burrows in arid regions of the Americas (Nguyen et al. 2013), produce arthroconidia. When arthroconidia are inhaled by a mammal, they give rise to round cells—spherules—in which growth and mitosis occur to eventually fill the enlarging mother cell with hundreds of endospores. On maturity, the spherule ruptures and releases the endospores into host tissue. Newly released endospores repeat the growth cycle locally or disseminate hematogenously. Phylogenetic analysis shows that the fungus evolved in North America before that continent's geologic connection with South America 2.5–3.5 million years ago (Fisher et al. 2001), and that the genetically homogeneous South

American strains, derived from populations in present-day Texas, reached their current distribution 8940–134,000 years ago, possibly with the migration of humans from North into South America (Fisher et al. 2001). Another unique member of the *Ajellomycetaceae*, *L. loboi*, shares an important feature with a distant member of its phylum Ascomycota, *Pneumocystis jirovecii*. They are the only human fungal pathogens not grown in culture (Vilela et al. 2009). *Lacazia*, a sister genus to *Paracoccidioides*, has been found only in subcutaneous infections of previously healthy humans in the Amazon basin and in dolphins of the Atlantic ocean and of the Amazon (Vilela et al. 2009; Theodoro et al. 2012), and (analogous to *Mycobacterium leprae*) can be amplified only in a mouse model. In contrast, *Paracoccidioides* spp., prevalent fungal pathogens that cause pneumonia, systemic disease of the monocyte–macrophage system, and destructive lesions of skin and oral mucous membranes (Fortes et al. 2011; Queiroz-Telles and Escuissato 2011; Marques 2012, 2013), occur in Central and South America mainly outside the Amazon region (Theodoro et al. 2012). *Paracoccidioides* causes clinical disease in many more men than women, in a ratio of 13:1, possibly because estrogen blocks the conversion of inhaled arthroconidia to the tissue-invasive yeast form (Shankar et al. 2011). This fungus frequently infects armadillos, and its virulence factors such as the immunodominant adhesin glycoprotein 43 (Puccia et al. 1999, 2008, 2011; Fortes et al. 2011) may have evolved in its coevolution with these ancient mammals (Bagagli et al. 2006; Richini-Pereira et al. 2009).

Some members of the *Onygenales* infect healthy humans, but very rarely cause invasive disease. Dermatophytes of the genus *Arthrodermataceae* specialize in degrading keratins, important structural proteins in vertebrate skin whose extensive cross-linking by disulfide bonds makes them proteinase resistant (Gradišar et al. 2005). With the evolution of animals with substantial keratin appendages such as scales, feathers, nails, and hair, specializing in use of these nutrient sources must have been a worthwhile niche for the presumed ancestor of

J.R. Köhler et al.

the dermatophytes (Weitzman and Summerbell 1995).

Reptilian amniotes had evolved keratin, so that the basis for this specialization was available as a fungal substrate 300 million years ago (Eckhart et al. 2008). Recent analysis of mitochondrial genomes of six dermatophytes comprising the three morphologically defined anamorphic genera, *Trichophyton*, *Microsporum*, and *Epidermophyton*, together with 29 other members of Ascomycota, confirmed that the dermatophytes are descended from a common ancestor (are monophyletic) and separated from other fungi only 32–50 million years ago (Wu et al. 2009). This study also found that classification of species by molecular characters frequently contradicted the traditional classifications based on morphologic traits (Wu et al. 2009), emphasizing the fact that fungal morphology is enormously flexible and convergent evolution is common, so that only limited predictions of their physiology can be made from the appearance of fungal structures. In contrast, molecular phylogenetic studies have confirmed the grouping of dermatophytes by their ecologic and clinical features, including the distinct sites of the human body they tend to infect (Ca-farchia et al. 2013).

Anthropophilic dermatophytes travel over the world with their human hosts. Although many species are endemic to specific geographic regions, some species occur worldwide. Their prevalence varies with the lifestyle and socioeconomic conditions of their human hosts, and is undergoing continuous epidemiologic changes (Ameen 2010). Unlike their thermal dimorphic Onygenales relatives, they can reach a new host by person-to-person transmission, and their access to new human substrate is made even easier by the ability of their arthroconidia to persist for years in fomites (Weitzman and Summerbell 1995). Anthropophilic dermatophytes, relying on person-to-person access to new hosts, can forgo long-distance travel via airborne conidia, which their geophilic relatives produce in larger profusion to reach distant deposits of animal keratin (Weitzman and Summerbell 1995).

Coevolution with their animal hosts presumably allowed dermatophytes to evade or

resist the evolutionarily more ancient innate immune mechanisms, so that today their control requires adaptive cellular immunity (Dahl 1993). By restricting their usual habitat to the most superficial keratinized layer of the skin and its appendages, dermatophytes reduce their contact with immune cells. Human-specific dermatophytes (anthropophiles) are able to down-regulate host inflammation to establish a chronic infection (Blake et al. 1991; Shiraki et al. 2006). In contrast, soil-dwelling (geophilic) and animal-specific (zoophilic) dermatophytes are unable to manipulate host immunity and are eliminated by a vigorous inflammatory response, illustrating the principle that a well-adapted parasite carefully calibrates its virulence.

Basidiomycota

Cryptococci

Basidiomycete yeast with a worldwide distribution, cryptococci, in the past century, infected humans only rarely (Molez 1998). In the 1950s, increasing numbers of cryptococcal meningo-encephalitis were reported from central Africa, viewed in retrospect as sentinels of the emergence of AIDS around the Congo River (Molez 1998). AIDS is still the setting in which the vast majority of cryptococcosis occurs (Mitchell and Perfect 1995; Pukkila-Worley and Mylonakis 2008); for 2006, 957,900 cases of cryptococcal meningitis associated with AIDS were estimated, resulting in 624,700 deaths (Park et al. 2009). Furthermore, recent outbreaks of *Cryptococcus gattii* infections with significant mortality on Vancouver Island and the northwestern United States raise the concern that the fungus may be evolving to become virulent for healthy humans (Fraser et al. 2005; Byrnes et al. 2010; Springer et al. 2012).

The pathogenic cryptococci, *C. neoformans* and its sister species *C. gattii*, enter a human by inhalation of infectious cells: dried yeast or possibly basidiospores, the products of meiosis after mating. When these small airborne cells, <5 µm in size, are inhaled, pneumonia can ensue in susceptible hosts, and on gaining access to

the bloodstream, the yeast can disseminate to all organs. *Cryptococcus* has a special predilection for the CNS and causes subacute meningoencephalitis, in which high intracranial pressure plays an especially deleterious role, which, if left untreated, is lethal. The fungus can persist for years in the lung or in sites of previous dissemination and reactivate only on weakened immune surveillance.

C. gattii causes invasive disease and death in wild and domestic land and ocean mammals and birds and pneumonia and meningoencephalitis in previously healthy people (Krockenberger et al. 2002; Miller et al. 2002; Raso et al. 2004; Santos et al. 2008; McGill et al. 2009). A wealth of analyses of pathogenic cryptococci over the past several decades have shown that *C. gattii* in many settings is a more common primary pathogen than *C. neoformans* (Speed and Dunt 1995; Springer et al. 2012), but new reports continue to blur the boundaries between primary and opportunistic pathogens among the *Cryptococcus* species (Chen et al. 2008; Chau et al. 2010; Choi et al. 2010), illustrating the limited utility of these terms (Casadevall and Pirofski 2001). A recent report has indicated that the presence of anti-GM-CSF autoantibody is a risk factor for *C. gattii* CNS infection in otherwise healthy individuals (Saijo et al. 2014). It suggests that patients with cryptococcal CNS infection considered “immunocompetent” may carry immune defects that can yet not be identified by routine immunological screening. An important condition for a clearer understanding of ecology, epidemiology, and evolution of cryptococci is that more resources for microbiologic analysis become available in poor countries. Cultures of blood and cerebrospinal fluid currently cannot be performed in medical settings in which most patients with cryptococcosis will seek help.

The separation of the sister species *C. gattii* and *C. neoformans* is thought to have occurred 45 million years ago (Simwami et al. 2011). Molecular typing recapitulates traditional serologic classification of varieties into *C. neoformans* var. *grubii* (serotype A, VNI, VNII, VNB) and var. *neoformans* (serotype D, VNIV) and their hybrids (serotype AD, VNIII), as well as *C. gattii*

(serotypes B and C) (Mitchell and Perfect 1995; Meyer et al. 2003). *C. gattii* has been subdivided into molecular varieties or cryptic species VGI through VGIV, among which further subdivisions correlate with geographic location and virulence (Kidd et al. 2004; Fraser et al. 2005; Byrnes et al. 2010).

In the environment, *C. gattii* has been isolated from numerous tree species in tropical and more recently in temperate regions, particularly from sites of wood decay and insect consumption of vegetable matter (Ellis and Pfeiffer 1990; Fortes et al. 2001; Kidd et al. 2003); although it was first found on *Eucalyptus* trees and believed to follow their worldwide distribution (Ellis and Pfeiffer 1990), its isolation from the midst of pristine Amazon rainforest revises the interpretation of its origin (Fortes et al. 2001). Environmental sampling on Vancouver Island to define the reservoirs of the outbreak that began in 1999 found more than 10 species of trees to yield *C. gattii*, which comprised ~10% of the trees sampled (Kidd et al. 2007), and it survives for at least 1 yr in fresh- and seawater, suggesting it could spread with ocean currents (Kidd et al. 2007).

Supporting the idea of an origin of pathogenic *C. gattii* in South America is the uniquely symmetric distribution of mating types in haploid isolates from that continent. Although clinical and environmental isolates of the α mating type vastly predominate in other parts of the world, and isolates of the α type are almost never encountered, the South American distribution is 0.8 α –1.0 α (Hagen et al. 2013).

C. neoformans var. *grubii*, which is responsible for ~95% of cryptococcal infections worldwide and 98% of infections among AIDS patients (Simwami et al. 2011), is genotypically most diverse in southern Africa; and a collection of strains from Botswana contained 12% isolates of the α mating type, which in other sites is extremely rare (Litvinseva and Mitchell 2012). In that study, distinct genotypes were thought to have diverged 5000 years ago; the investigators speculate that pigeons contributed to the worldwide spread of one specific strain of *C. neoformans* var. *grubii* (Litvinseva and Mitchell 2012) because *C. neoformans* is

J.R. Köhler et al.

enriched in bird droppings and has been shown to mate on pigeon droppings (Nielsen et al. 2007). A study from Thailand supports the concept of an origin of *C. neoformans* var. *grubii* in Africa (Simwami et al. 2011).

Like *C. neoformans*, with which it shares virulence traits, and unlike nonpathogenic cryptococcal species (Findley et al. 2009; Araujo Gde et al. 2012), *C. gattii* possesses a thick polysaccharide capsule composed of glucurono- and galactoxylomannan, which, analogous to encapsulated bacteria like *Streptococcus pneumoniae*, blocks phagocytosis of the organism unless it is opsonized (Casadevall et al. 1998). Presumably, the dense capsule also provides some protection against hydrolytic enzymes exocytosed by neutrophils. Capsular glucuronoxylomannan and other macromolecules, as well as virulence-associated proteins, are carried through the cell wall in an amazing process involving vesicles that originated from the late endosome (multivesicular body) (Rodrigues et al. 2007, 2008). This process, first discovered in *Cryptococcus*, subsequently was found in other pathogenic fungi like *Paracoccidioides* (Vallejo et al. 2011, 2012), and in the model yeast *Saccharomyces cerevisiae* (Oliveira et al. 2010). Like other fungi, cryptococci can diversify morphologically in the host, producing a population of Titan cells, giant cells of up to 50 μm in diameter (Okagaki et al. 2010; Zaragoza et al. 2010) created by DNA replication and growth not followed by mitosis (Okagaki et al. 2010). Other antiphagocytic mechanisms, independent of the capsule, are controlled by two GATA transcription factors and are likely to involve coordinated regulation of numerous physiologic processes because hundreds of genes are differentially expressed in their mutants (Liu et al. 2008; Chun et al. 2011).

Once ingested by alveolar macrophages not activated by T lymphocytes, the fungus can survive and proliferate, and it is thought that some cryptococci may enter the brain within macrophages crossing the blood–brain barrier (a Trojan horse-like mechanism), whereas free yeast in the bloodstream cross the blood–brain barrier by transcytosis (Casadevall 2010; Shi et al. 2010). To survive in phagocytes, it buffers

oxidative and nitrosative stress with melanin deposited in its cell wall (Wang et al. 1995), and with neutralizing enzymes like superoxide dismutase, glutathione reductase, and thioredoxins (Kronstad et al. 2011). Both in macrophages and in the free-living amoeba *Acanthamoeba castellanii*, it disrupts phagolysosome membranes to prevent acidification of this compartment and to dilute the lytic enzymes it contains; it also sheds large amounts of capsular polysaccharide to fill the phagocytic cells' cytoplasm (Steenbergen et al. 2001; Tucker and Casadevall 2002). Virulent strains not only resist killing but proliferate in the amoebae (Steenbergen et al. 2001) so that the predator finds itself prey.

How did some cryptococci evolve from survivors of amoeba attacks to occasional parasites of humans? A recent study examined evolutionary relationships and virulence traits of *C. gattii* and *C. neoformans* strains, and related fungi belonging to the order Tremellales, which do not infect mammals (Findley et al. 2009). Other Tremellales parasitize fungi by attaching to the host fungus with a specialized hypha and accessing its cytoplasm through a newly created pore (Zugmaier et al. 1994; Millanes et al. 2014). An evolutionary trajectory from parasitizing another live fungus to parasitizing an animal host may have proceeded through association with and parasitism of insects, and through selection in soils by amoebae for traits that can also function in mammalian virulence. The human-pathogenic *Cryptococcus* species killed larvae of the wax moth *Galleria mellonella* in one study, whereas their Tremellales relatives partially or completely lacked virulence in this insect model (Findley et al. 2009). Of note, insect cellular immune defenses consist of amoeboid phagocytes (Williams 2007; Browne et al. 2013), so that resistance to free-living amoebae may transfer well to resistance to insect plasmatocytes. However, in nature, cryptococci so far have only been found in association with insect frass (the excreta of plant-eating insects), not as insect parasites (unlike, e.g., the entomophthorales and the ascomycetous *Cordyceps* spp.). Further ecologic research will have to show whether insect parasitic cryptococcal relatives can be

identified, perhaps in their ancestral tropical forest habitats. The role of its versatile mating system in virulence evolution of *Cryptococcus* will be discussed in the dedicated articles in this collection.

FUNGI THAT INFECT IMMUNOCOMPROMISED HUMANS

A sufficiently immunocompromised host can be infected by hundreds of environmental fungal species that grow at human core temperatures. However, a predictable set of actors is known to cause the most common invasive infections in immunocompromised individuals, and they will be discussed next according to their phylogenetic affiliations, which predict important parameters of their physiology.

Ascomycota

Candida

Candida species are a polyphyletic group of the order Saccharomycotina, which live as commensals on mammalian mucous membranes, particularly of the gastrointestinal tract (Wrobel et al. 2008). They have rarely been found in the soil (Marples and Di Menna 1952; Skinner and Fletcher 1960). A limited number of species are commonly associated with humans as colonizers and opportunistic pathogens: *C. albicans*, *Candida glabrata*, *Candida parapsilosis*, *Candida tropicalis*, *Candida lusitanae*, and *Candida krusei*. Of these, *C. albicans* is the most commonly isolated human commensal and pathogen (Odds 1988; Kam and Xu 2002; Krcmery and Barnes 2002). The frequency of colonization with nonalbicans *Candida* species shifts according to age of the host, with *C. parapsilosis* being more prevalent among children and *C. glabrata* among older adults (Soll 2002).

Invasive candidiasis takes many forms, depending on the setting in which a host became susceptible to this opportunist, and depletion of normally competing bacterial flora by antibiotics often plays a role. For example, in cancer patients receiving chemotherapy, candidiasis is often caused by fungal transmigration through

The Spectrum of Fungi That Infects Humans

disrupted intestinal epithelium, and the organs with the heaviest fungal burden are those connected by the portal circulation—liver and spleen—but not the brain. In contrast, premature newborns with candidiasis often have abscesses in the brain, because their blood–brain barrier may be immature and because their brain receives such a large fraction of cardiac blood output.

C. albicans is the human companion fungus par excellence. Many healthy humans carry *C. albicans*, which harmlessly colonizes mucous membranes to high numbers at different anatomic sites of a single individual (Odds 1984, 1988; Soll et al. 1991). It has evolved to flourish in a wide range of environmental conditions: high pH in the intestine versus low pH in the vagina, feast and famine nutritional conditions in the gastrointestinal tract according to its host's mealtimes versus steady glycogen supply in the vagina, and aerobic conditions on oral surfaces versus anaerobic conditions in the intestine. Unlike the dermatophytes, in which different clades inhabit different anatomic sites, *C. albicans* biotypes are generalists and each can adapt to all sites (Odds 1984).

Not only can *C. albicans* survive and thrive in highly disparate microenvironments of the human host, but as part of the normal flora it has also honed its ability to avoid triggering human immune defenses, for instance, by covering its main structural cell wall component, $\beta(1,3)$ -D-glucan, with glycoproteins to avoid engaging the macrophage dectin-1 receptor (Wheeler and Fink 2006).

Analysis of its populations, which comprise at least five clades, suggests that *C. albicans* migrated throughout the world with its human hosts (Lott et al. 2005). It is unknown whether the fungus accompanied the mammalian lineages that led to primates throughout evolution, or whether it made the jump to mammalian commensal at a more recent time before emergence of modern humans (Lott et al. 2005).

Perhaps in its long cohabitation with the human, which as to host numbers proved to be a winning strategy, *C. albicans* became so closely hostadapted that major genomic rearrangements, as facilitated by sex and meiotic recom-

J.R. Köhler et al.

bination, became less advantageous (Goddard et al. 2005). In a stable and benign environment, such as the equilibrium between thriving hosts and commensals, the expenditure of time and energy required for sexual reproduction may cost more than it benefits. *C. albicans* appears to consist of stable clonal diploid populations (Bougnoux et al. 2008), and its mating and parasexual cycle seem to be such rare events that their discoveries were sensational paradigm shifts in the community of *Candida* researchers (Hull and Johnson 1999; Magee and Magee 2000; Miller and Johnson 2002; Alby et al. 2009; Hickman et al. 2013). Although the parasexual cycle occurs infrequently, it can generate progeny with extensive genetic diversity by shuffled combinations of eight chromosomes or by recombination between homologous chromosomes mediated by the conserved Spo11 protein integral to meiotic recombination (Forche et al. 2008). More discoveries are sure to follow on the role of mating and various ploidy states in *Candida* ecology today, and virulence associated with the mating loci (Lockhart et al. 2005; Wu et al. 2007) may become more important as *Candida* habitat changes with current changes in human epidemiology.

In the millions of years of coevolution, human predecessors and humans survived only briefly with diseases during which *C. albicans* could significantly invade the host. Invasive candidiasis was very rare (Browne 1954; Zimmerman 1955). *C. albicans* is an efficient invasive pathogen, causing mucous membrane infections in individuals with ineffective adaptive cellular immunity and fatal disseminated infections in patients lacking functional innate immune cells, neutrophils. For a microorganism that relies on person-to-person transmission and does not have a significant soil reservoir (or mode of locomotion back from the soil to its primary host), the benefit of multiplying to the point of host death seems mysterious. One idea might be that the *Candida* pathogenic repertoire evolved to take advantage of a constant large number of temporarily susceptible humans, infants, who develop oral candidiasis because of immature adaptive immunity. Human infants' propensity to widely distribute their

oral secretions (to slobber) may allow infecting *Candida* to launch its increased numbers from infected mucous membranes of these hosts and to colonize more of their contacts. If this is the case, *Candida* virulence factors evolved to efficiently infect mucous membranes, and invasive disease is for it an accidental dead end.

The pathogenic repertoire of *Candida* comprises all features needed for a human fungal pathogen. It is able to grow well at human febrile temperatures of 39°C–40°C. It can penetrate host tissues with hyphal cells and it has multiple adhesion molecules to facilitate the drilling action of the hyphal tip (Staab et al. 1999; Sundstrom 2002; Hoyer et al. 2008; Liu and Filler 2011). Its hyphae constitutively produce yeast, constantly diversifying the population with more mobile cells (Shen et al. 2008). An array of lytic enzymes suitable for digesting human macromolecules is induced during tissue invasion (Ghannoum 2000; Staib et al. 2000; Calderone and Fonzi 2001; Schaller et al. 2005; Albrecht et al. 2006; Trofa et al. 2011), and transporters are regulated coordinately to absorb the released monomers into the fungal cell, as reviewed by (Morschhäuser 2011), and to mobilize micronutrients from the host (Weissman et al. 2008; Citiulo et al. 2012; Noble 2013). Unbiased genetic screening yielded novel virulence factors like glucosylceramide synthesis (Noble et al. 2010) and is expected to reveal new aspects of *Candida* invasive lifestyle (Holland and Summers 2008).

Resistance of *Candida* to the human immune system is limited. Features increasing its virulence were likely selected against in its long coevolution with us as a commensal. Yet today we are saving, and improving the quality of, human lives that are vulnerable to this opportunist, and any line of investigation that might result in better control of invasive candidiasis has the potential worldwide to save hundreds of thousands of lives every year (Pfaller and Diekema 2007).

Aspergillus

Aspergillus fumigatus, the species responsible for ~90% of invasive aspergillosis (Schmitt

et al. 1990), causes relentless pneumonia, sinusitis progressing through tissue planes, and brain abscesses in neutropenic patients and those with phagocyte defects like chronic granulomatous disease. It is feared in immunosuppressed individuals because its susceptibility to antifungals is limited. This organism also causes disease at another extreme of immune function: allergic reactions.

Normal human innate immunity controls *A. fumigatus*, a versatile thermophilic plant saprobe. It is well adapted to the high temperatures that occur during bacterial decomposition of dead plants and tolerates thermophilic bacteria's optimum of around 55°C (Latgé 1999). At human febrile temperatures of 39°C–40°C, it grows well. Its secretion of a very broad and redundant range of hydrolases (Kothary et al. 1984; Kwon-Chung and Sugui 2013), evolved in the tumultuous competition for nutrients among the microbiome of decaying plants, allows it to easily access human tissues as sources of nutrition (Abad et al. 2010).

The mycelia of *A. fumigatus* give rise to conidiophores, specialized hyphae that produce small (~3 µm) airborne dispersal cells, conidia. Their small size allows them to remain airborne for long periods and, incidentally, to enter human alveoli. On average, a person is estimated to inhale several hundred *A. fumigatus* conidia each day (Hospenthal et al. 1998); with exposure to grass cuttings, leaf litter, or compost, this number may be orders of magnitude higher (Mullins et al. 1976; Poole and Wong 2013). Conidia are coated with hydrophobic proteins (Thau et al. 1994; Paris et al. 2003) and with the chemoprotectant melanin (Pihet et al. 2009) to withstand harsh environmental stressors like freezing, sunlight, and desiccation (Kwon-Chung and Sugui 2013). They are recognized by innate immune cells of a human host only when they begin to germinate to produce a growing hypha (Levitz and Diamond 1985; Aimagani et al. 2009) after landing in an air-filled space like a paranasal sinus or lung. *Aspergillus* conidia can give rise to allergic lung disease, and the fungus can colonize the bronchiectasis of cystic fibrosis or chronic obstructive lung disease; but lethal invasive infection

occurs when there is a dearth of functional neutrophils and macrophages to control germinating conidia. Hyphae proliferate in lung parenchyma or sinuses, and hyphal fragments occasionally are carried to highly perfused organs like the brain in which they initiate foci of metastatic infection. On encountering blood vessels, *Aspergillus* hyphae tend to enter and follow their course, clogging the vessel and causing infarction of downstream tissue. This angioinvasive behavior may be attributable to thigmotropism, the ability to sense and follow contours, which aspergilli have in common with fungi of diverse phyla (Perera et al. 1997; Bowen et al. 2007).

Pathogenicity studies of *A. fumigatus* highlight general themes of innate immunity (Morton et al. 2012). For example, specialized alveolar epithelial cells, type II pneumocytes, phagocytose and kill conidia of *A. fumigatus* in their lysosomes, unless the conidia manage to germinate before their death (Wasylnka and Moore 2003). Neutrophils are attracted by the chemokines secreted by resident alveolar macrophages encountering *A. fumigatus* during their patrols of the alveolar space. Whereas small hyphae are engulfed and killed by neutrophils, larger hyphae are killed by their extracellular release of reactive oxygen and nitrogen species and antimicrobial peptides, in dependence on the pathogen recognition receptors (PRRs) TLR2 and 4, and dectin-1 (Netea et al. 2003; Kennedy et al. 2007; Werner et al. 2009). Neutrophil extracellular traps (NETs), critical to control polymicrobial infection in appendicitis (Brinkmann et al. 2004), nutritionally constrain *A. fumigatus* hyphae (McCormick et al. 2010) and down-regulate excessive inflammation elicited by their presence (Rohm et al. 2014).

Non-Fumigatus Aspergilli, Fusarium, Pseudoallescheria, and Other Opportunistic Ascomycetous Fungal Pathogens

Profound immunosuppression, for instance, during prolonged neutropenia, graft-versus-host disease, or severe rejection episodes of transplanted solid organs, permits invasive disease of many other types of environmental fila-

J.R. Köhler et al.

mentous ascomycetes. Among them are the non-*fumigatus* species of *Aspergillus* (Torres et al. 2003), *Fusarium solanum*, *Fusarium oxysporum*, and other *Fusarium* spp., and their teleomorph *Nectria* spp. (Nucci and Anaissie 2007), *Pseudoallescheria boydii*, and its *Scedosporium anamorphs* (Quan and Spellberg 2010). Infections with these fungi are often lethal because the hosts they usually infect are incapable of mounting an effective immune response, and because they tend to be more resistant to currently available antifungals.

Fusarium spp. have evolved to infect plants, and the genomes of certain lineages contain one or more entire chromosomes encoding plant pathogenicity factors, which may be horizontally transferred within the genus (Ma et al. 2010). Encoded on pathogenicity chromosomes, and other genomic clusters reminiscent of bacterial pathogenicity islands, are secreted hydrolytic enzymes and signaling molecules expressed during early plant infection (Ma et al. 2010; Rep and Kistler 2010). Whether possession of a pathogenicity chromosome corresponds to virulence for humans, for instance, in the *F. oxysporum* lineage identified to have caused 70% of invasive fusariosis in a San Antonio hospital (O'Donnell et al. 2004), is not yet known. In *Fusarium*, morphogenesis contributes to virulence for humans, because yeast-like cells are produced from hyphae in the host, and typically spread widely through the bloodstream to cause numerous foci of infection in the skin.

Pigmented filamentous ascomycetes, also called dematiaceous fungi, infect immunocompromised and, rarely, immunocompetent individuals to cause phaeohyphomycosis. Their pathobiology is diverse including brain abscesses (*Cladophiala bantianum*, *Ramichloridium* spp., and *Dactylaria gallopava*), keratitis, sinus or soft tissue infections, ulcers, and cysts (*Exophiala jeanselmei*, *Exophiala dermatitidis*, *Curvularia* spp., *Bipolaris*, or *Alternaria* spp.). When directly inoculated into tissue, these fungi can cause devastating disease, as emphasized in a tragic outbreak caused by contaminated injectable steroids (Smith et al. 2013).

Another epidemiologically, biologically, and profoundly important ascomycete, *P. jiro-*

vecii, is so distinctive in its biology, that only a reference to the related article in this collection can be made (Gigliotti et al. 2014).

Basidiomycota

In addition to the *Cryptococcus* spp. discussed above, other Basidiomycota growing predominantly as yeast in the host, *Malassezia furfur*, *Trichosporon asahii*, and other members of human skin flora, are opportunists in patients with venous catheters and in immunosuppressed patients.

Mucorales

Profoundly immunocompromised patients at risk for the environmental fungi described above are also at risk for Mucorales infections. Additionally, Mucorales cause severe disease in diabetic patients especially at times of uncontrolled blood glucose, and patients with elevated serum iron (e.g., attributed to hemosiderosis), as described in the excellent series by Roden et al. (2005). Like the aspergilli, these fungi cause disease where airborne dispersal cells, sporangiospores, enter air-filled spaces (i.e., in sinuses and lungs). Infections with Mucorales progress rapidly as the fungus grows fast and relentlessly through tissue planes and bone, penetrating the eye and the brain when originating in sinuses, and causing widespread infarction because of its angioinvasive behavior. Depending on the underlying condition, such infections are often fatal (Roden et al. 2005), especially because these fungi tend toward resistance to current antifungals.

The Mucorales are an ubiquitous, ancient group mostly of saprotrophs, which, unlike the Ascomycota and Basidiomycota (the Dikarya), do not manifest sophisticated adaptations to diverse substrates, but rapidly use up available sugars and then move on via their sporangiospores (Hoffmann et al. 2013). Their evolutionary distance from the phyla of other human parasitic fungi is apparent in the structure of their hyphae, which are fragile, thin walled, and lack septa, and in the composition of their cell walls in which chitin and chitosan

play the structural roles that glucans fulfill in the Dikarya (Dijksterhuis and Samson 2006). Some species like *Mucor circillienoides* are dimorphic and form yeast in the host (Dizbay et al. 2009; Khan et al. 2009; Lee et al. 2013), contributing to dissemination.

Among the *Rhizopus* spp., *Rhizopus oryzae* is responsible for ~70% of human disease (Spellberg et al. 2005). Several of its virulence factors are areas of active investigation: its iron scavenging from the host (Fu et al. 2004; Ibrahim et al. 2008, 2010; Spellberg et al. 2009; Ibrahim 2011) and its ability to bind GRP78/BiP, a Kar2 homolog, on the surface of endothelial cells with its spore coat protein homolog Coth3 (Liu et al. 2010; Gebremariam et al. 2014). Like a broad phylogenetic variety of fungi, Mucorales are angioinvasive. Thigmotropism, the proclivity to sense and follow curvatures of a surface, has been shown for one species, *Mucor mucedo* (Perera et al. 1997), and is likely to be common to all. Much remains to be learned about the biology of these evolutionarily old fungi and historically new human pathogens.

CONCLUSION

Immunologically intact humans manifest robust defenses against fungal diseases. Recently, human social evolution produced scientific medicine, whose progress has rendered a large population susceptible to infections with fungi not considered human pathogens as recently as a hundred years ago. Human social evolution may have reached a stage where prioritization of the resources needed for understanding fungal biology and for successful development of multiple new classes of antifungals is possible, as the example of AIDS shows. Effective antiretroviral therapy was developed with a massive research effort less than two decades ago, and if human society evolves to a point of valuing all human lives equally, near elimination of AIDS is as feasible as the quasi elimination of HIV mother-to-child transmission has been in wealthy countries (Siegfried et al. 2011; Cohen et al. 2013; Nicol et al. 2013). Unlike with HIV, which has no reservoir in nature, humans will always have to cope with fungal infections, because poten-

The Spectrum of Fungi That Infects Humans

tial fungal pathogens are part of our normal flora and of soil, water, and air. But building on the tremendous efforts and findings of the past decades, new low-toxicity therapies and novel preventative measures of fungal infections can be within our reach.

ACKNOWLEDGMENTS

J.R.K. thanks Alison Clapp, Library Director of Boston Children's Hospital, for ever-gracious and lightning-quick procurement of innumerable articles.

REFERENCES

*Reference is also in this collection.

- Abad A, Fernandez-Molina JV, Bikandi J, Ramirez A, Margareto J, Sendino J, Hernando FL, Ponton J, Garaizar J, Rementeria A. 2010. What makes *Aspergillus fumigatus* a successful pathogen? Genes and molecules involved in invasive aspergillosis. *Rev Iberoam Micol* **27**: 155–182.
- Aimanianda V, Bayry J, Bozza S, Knemeyer O, Perruccio K, Elluru SR, Clavaud C, Paris S, Brakhage AA, Kaveri SV, et al. 2009. Surface hydrophobin prevents immune recognition of airborne fungal spores. *Nature* **460**: 1117–1121.
- Albrecht A, Felk A, Pichova I, Naglik JR, Schaller M, de Groot P, Maccallum D, Odds FC, Schafer W, Klis F, et al. 2006. Glycosylphosphatidylinositol-anchored proteases of *Candida albicans* target proteins necessary for both cellular processes and host–pathogen interactions. *J Biol Chem* **281**: 688–694.
- Alby K, Schaefer D, Bennett RJ. 2009. Homothallic and heterothallic mating in the opportunistic pathogen *Candida albicans*. *Nature* **460**: 890–893.
- Ameen M. 2010. Epidemiology of superficial fungal infections. *Clin Dermatol* **28**: 197–201.
- Araujo Gde S, Fonseca FL, Pontes B, Torres A, Cordero RJ, Zancoppe-Oliveira RM, Casadevall A, Viana NB, Nimrichter L, Rodrigues ML, et al. 2012. Capsules from pathogenic and non-pathogenic *Cryptococcus* spp. manifest significant differences in structure and ability to protect against phagocytic cells. *PLoS ONE* **7**: e29561.
- Argimon S, Fanning S, Blankenship JR, Mitchell AP. 2011. Interaction between the *Candida albicans* high-osmolarity glycerol (HOG) pathway and the response to human β -defensins 2 and 3. *Eukaryot Cell* **10**: 272–275.
- Aulakh HK, Aulakh KS, Troy GC. 2012. Feline histoplasmosis: A retrospective study of 22 cases (1986–2009). *J Am Anim Hosp Assoc* **48**: 182–187.
- Bagagli E, Bosco SM, Theodoro RC, Franco M. 2006. Phylogenetic and evolutionary aspects of *Paracoccidioides brasiliensis* reveal a long coexistence with animal hosts that explain several biological features of the pathogen. *Infect Genet Evol* **6**: 344–351.

J.R. Köhler et al.

- Bagagli E, Theodoro RC, Bosco SM, McEwen JG. 2008. *Paracoccidioides brasiliensis*: Phylogenetic and ecological aspects. *Mycopathologia* **165**: 197–207.
- Bastidas RJ, Heitman J. 2009. Trimorphic stepping stones pave the way to fungal virulence. *Proc Natl Acad Sci* **106**: 351–352.
- Bauder B, Kubber-Heiss A, Steineck T, Kuttin ES, Kaufman L. 2000. Granulomatous skin lesions due to histoplasmosis in a badger (*Meles meles*) in Austria. *Med Mycol* **38**: 249–253.
- Bergman A, Casadevall A. 2010. Mammalian endothermy optimally restricts fungi and metabolic costs. *mBio* **1**: e00212-10.
- Beyhan S, Gutierrez M, Voorhies M, Sil A. 2013. A temperature-responsive network links cell shape and virulence traits in a primary fungal pathogen. *PLoS Biol* **11**: e1001614.
- Bittencourt AL. 1988. Entomophthoromycosis. Review. *Med Cutan Ibero Lat Am* **16**: 93–100.
- Bittencourt AL, Marback R, Nossa LM. 2006. Mucocutaneous entomophthoromycosis acquired by conjunctival inoculation of the fungus. *Am J Trop Med Hyg* **75**: 936–938.
- Blake JS, Dahl MV, Herron MJ, Nelson RD. 1991. An immunoinhibitory cell wall glycoprotein (Mannan) from *Trichophyton rubrum*. *J Invest Dermatol* **96**: 657–661.
- Boelaert JR, de Locht M, Van Cutsem J, Kerrels V, Cantiniaux B, Verdonck A, Van Landuyt HW, Schneider YJ. 1993. Mucormycosis during deferoxamine therapy is a siderophore-mediated infection. In vitro and in vivo animal studies. *J Clin Invest* **91**: 1979–1986.
- Bohse ML, Woods JP. 2007. Expression and interstrain variability of the *YPS3* gene of *Histoplasma capsulatum*. *Eukaryot Cell* **6**: 609–615.
- Bougnoux ME, Pujol C, Diogo D, Bouchier C, Soll DR, d'Enfert C. 2008. Mating is rare within as well as between clades of the human pathogen *Candida albicans*. *Fungal Genet Biol* **45**: 221–231.
- Bowen AD, Davidson FA, Keatch R, Gadd GM. 2007. Induction of contour sensing in *Aspergillus niger* by stress and its relevance to fungal growth mechanics and hyphal tip structure. *Fungal Genet Biol* **44**: 484–491.
- Bradsher RW, Chapman SW, Pappas PG. 2003. Blastomycosis. *Infect Dis Clin North Am* **17**: 21–40.
- Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y, Zychlinsky A. 2004. Neutrophil extracellular traps kill bacteria. *Science* **303**: 1532–1535.
- Brömel C, Sykes JE. 2005. Histoplasmosis in dogs and cats. *Clin Tech Small Anim Pract* **20**: 227–232.
- Browne SG. 1954. Moniliasis following antibiotic therapy. *Lancet* **266**: 393–395.
- Browne N, Heelan M, Kavanagh K. 2013. An analysis of the structural and functional similarities of insect hemocytes and mammalian phagocytes. *Virulence* **4**: 597–603.
- Byrnes EJ III, Li W, Lewit Y, Ma H, Voelz K, Ren P, Carter DA, Chaturvedi V, Bildfell RJ, May RC, et al. 2010. Emergence and pathogenicity of highly virulent *Cryptococcus gattii* genotypes in the northwest United States. *PLoS Pathog* **6**: e1000850.
- Cafarchia C, Iatta R, Latrofa MS, Graser Y, Otranto D. 2013. Molecular epidemiology, phylogeny and evolution of dermatophytes. *Infect Genet Evol* **20**: 336–351.
- Cain CW, Lohse MB, Homann OR, Sil A, Johnson AD. 2012. A conserved transcriptional regulator governs fungal morphology in widely diverged species. *Genetics* **190**: 511–521.
- Calderone RA, Fonzi WA. 2001. Virulence factors of *Candida albicans*. *Trends Microbiol* **9**: 327–335.
- Casadevall A. 2005. Fungal virulence, vertebrate endothermy, and dinosaur extinction: Is there a connection? *Fungal Genet Biol* **42**: 98–106.
- Casadevall A. 2010. Cryptococci at the brain gate: Break and enter or use a Trojan horse? *J Clin Invest* **120**: 1389–1392.
- Casadevall A. 2012. Fungi and the rise of mammals. *PLoS Pathog* **8**: e1002808.
- Casadevall A, Pirofski L. 2001. Host–pathogen interactions: The attributes of virulence. *J Infect Dis* **184**: 337–344.
- Casadevall A, Pirofski LA. 2007. Accidental virulence, cryptic pathogenesis, Martians, lost hosts, and the pathogenicity of environmental microbes. *Eukaryot Cell* **6**: 2169–2174.
- Casadevall A, Cleare W, Feldmesser M, Glatman-Freedman A, Goldman DL, Kozel TR, Lendvai N, Mukherjee J, Pirofski LA, Rivera J, et al. 1998. Characterization of a murine monoclonal antibody to *Cryptococcus neoformans* polysaccharide that is a candidate for human therapeutic studies. *Antimicrob Agents Chemother* **42**: 1437–1446.
- Chau TT, Mai NH, Phu NH, Nghia HD, Chuong LV, Sinh DX, Duong VA, Diep PT, Campbell JI, Baker S, et al. 2010. A prospective descriptive study of cryptococcal meningitis in HIV uninfected patients in Vietnam—High prevalence of *Cryptococcus neoformans* var *grubii* in the absence of underlying disease. *BMC Infect Dis* **10**: 199.
- Chen J, Varma A, Diaz MR, Litvintseva AP, Wollenberg KK, Kwon-Chung KJ. 2008. *Cryptococcus neoformans* strains and infection in apparently immunocompetent patients, China. *Emerg Infect Dis* **14**: 755–762.
- Chen YL, Lehman VN, Lewit Y, Averette AF, Heitman J. 2013. Calcineurin governs thermotolerance and virulence of *Cryptococcus gattii*. *G3 (Bethesda)* **3**: 527–539.
- Choi YH, Ngamskulrungroj P, Varma A, Sionov E, Hwang SM, Carriconde F, Meyer W, Litvintseva AP, Lee WG, Shin JH, et al. 2010. Prevalence of the VN1c genotype of *Cryptococcus neoformans* in non-HIV-associated cryptococcosis in the Republic of Korea. *FEMS Yeast Res* **10**: 769–778.
- Chun CD, Brown JC, Madhani HD. 2011. A major role for capsule-independent phagocytosis-inhibitory mechanisms in mammalian infection by *Cryptococcus neoformans*. *Cell Host Microbe* **9**: 243–251.
- Citiulo F, Jacobsen ID, Miramon P, Schild L, Brunke S, Zipfel P, Brock M, Hube B, Wilson D. 2012. *Candida albicans* scavenges host zinc via Pra1 during endothelial invasion. *PLoS Pathog* **8**: e1002777.
- Cohen MS, Smith MK, Muessig KE, Hallett TB, Powers KA, Kashuba AD. 2013. Antiretroviral treatment of HIV-1 prevents transmission of HIV-1: Where do we go from here? *Lancet* **382**: 1515–1524.

- Comerio RM, Andorno AV, Botto EN. 2008. *Conidiobolus coronatus* isolation from a pest aphids of chives (*Allium schoenoprasum* L.). *Rev Iberoam Micol* **25**: 193–195.
- Cramer RA Jr, Perfect BZ, Pinchai N, Park S, Perlin DS, Asfaw YG, Heitman J, Perfect JR, Steinbach WJ. 2008. Calcineurin target CrzA regulates conidial germination, hyphal growth, and pathogenesis of *Aspergillus fumigatus*. *Eukaryot Cell* **7**: 1085–1097.
- Dahl MV. 1993. Suppression of immunity and inflammation by products produced by dermatophytes. *J Am Acad Dermatol* **28**: S19–S23.
- Dijksterhuis J, Samson RA. 2006. Zygomycetes. In *Food spoilage microorganisms* (ed. Blackburn CW), pp. 415–436. CRC, Boca Raton, FL.
- Dizbay M, Adisen E, Kustimur S, Sari N, Cengiz B, Yalcin B, Kalkanci A, Gonullu, Sugita T. 2009. Fungemia and cutaneous zygomycosis due to *Mucor circinelloides* in an intensive care unit patient: Case report and review of literature. *Jpn J Infect Dis* **62**: 146–148.
- Echeteu CO, Ononogbu IC. 1982. Extracellular lipase and proteinase of *Basidiobolus haptosporus*: Possible role in subcutaneous mycosis. *Mycopathologia* **80**: 171–177.
- Eckhart L, Valle LD, Jaeger K, Ballaun C, Szabo S, Nardi A, Buchberger M, Hermann M, Alibardi L, Tschachler E. 2008. Identification of reptilian genes encoding hair keratin-like proteins suggests a new scenario for the evolutionary origin of hair. *Proc Natl Acad Sci* **105**: 18419–18423.
- Edwards JA, Alore EA, Rappleye CA. 2011. The yeast–phase virulence requirement for α -glucan synthase differs among *Histoplasma capsulatum* chemotypes. *Eukaryot Cell* **10**: 87–97.
- Edwards JA, Chen C, Kemski MM, Hu J, Mitchell TK, Rappleye CA. 2013. *Histoplasma* yeast and mycelial transcriptomes reveal pathogenic-phase and lineage-specific gene expression profiles. *BMC Genomics* **14**: 695.
- Ellis DH, Pfeiffer TJ. 1990. Natural habitat of *Cryptococcus neoformans* var. *gattii*. *J Clin Microbiol* **28**: 1642–1644.
- Ernst JF, Pla J. 2011. Signaling the glycoshield: Maintenance of the *Candida albicans* cell wall. *Int J Med Microbiol* **301**: 378–383.
- Findley K, Rodriguez-Carres M, Metin B, Kroiss J, Fonseca A, Vilgalys R, Heitman J. 2009. Phylogeny and phenotypic characterization of pathogenic *Cryptococcus* species and closely related saprobic taxa in the Tremellales. *Eukaryot Cell* **8**: 353–361.
- Fisher MC, Koenig GL, White TJ, San-Blas G, Negroni R, Alvarez IG, Wanke B, Taylor JW. 2001. Biogeographic range expansion into South America by *Coccidioides immitis* mirrors New World patterns of human migration. *Proc Natl Acad Sci* **98**: 4558–4562.
- Forche A, Alby K, Schaefer D, Johnson AD, Berman J, Bennett RJ. 2008. The parasexual cycle in *Candida albicans* provides an alternative pathway to meiosis for the formation of recombinant strains. *PLoS Biol* **6**: e110.
- Fortes ST, Lazera MS, Nishikawa MM, Macedo RC, Wanke B. 2001. First isolation of *Cryptococcus neoformans* var. *gattii* from a native jungle tree in the Brazilian Amazon rainforest. *Mycoses* **44**: 137–140.
- Fortes MR, Miot HA, Kurokawa CS, Marques ME, Marques SA. 2011. Immunology of paracoccidioidomycosis. *An Bras Dermatol* **86**: 516–524.
- Fraser JA, Giles SS, Wenink EC, Geuness-Boyer SG, Wright JR, Diezmann S, Allen A, Stajich JE, Dietrich FS, Perfect JR, et al. 2005. Same-sex mating and the origin of the Vancouver Island *Cryptococcus gattii* outbreak. *Nature* **437**: 1360–1364.
- Fu Y, Lee H, Collins M, Tsai HF, Spellberg B, Edwards JE Jr, Kwon-Chung KJ, Ibrahim AS. 2004. Cloning and functional characterization of the *Rhizopus oryzae* high affinity iron permease (*rFTR1*) gene. *FEMS Microbiol Lett* **235**: 169–176.
- Garcia-Solache MA, Casadevall A. 2010. Global warming will bring new fungal diseases for mammals. *mBio* **1**: e00061-10.
- Gebremariam T, Liu M, Luo G, Bruno V, Phan QT, Waring AJ, Edwards JE Jr, Filler SG, Yeaman MR, Ibrahim AS. 2014. CotH3 mediates fungal invasion of host cells during mucormycosis. *J Clin Invest* **124**: 237–250.
- Ghannoum MA. 2000. Potential role of phospholipases in virulence and fungal pathogenesis. *Clin Microbiol Rev* **13**: 122–143.
- * Gigliotti F, Limper AH, Wright T. 2014. *Pneumocystis*. *Cold Spring Harb Perspect Med*. doi: 10.1101/cshperspect.a019828.
- Goddard MR, Godfray HC, Burt A. 2005. Sex increases the efficacy of natural selection in experimental yeast populations. *Nature* **434**: 636–640.
- Gradisar H, Friedrich J, Krizaj I, Jerala R. 2005. Similarities and specificities of fungal keratinolytic proteases: Comparison of keratinases of *Paecilomyces marquandii* and *Doratomyces microsporus* to some known proteases. *Appl Environ Microbiol* **71**: 3420–3426.
- Gugnani HC. 1992. Entomophthoromycosis due to *Conidiobolus*. *Eur J Epidemiol* **8**: 391–396.
- Hagen F, Ceresini PC, Polacheck I, Ma H, van Nieuwerburgh F, Gabaldon T, Kagan S, Pursall ER, Hoogveld HL, van Iersel LJ, et al. 2013. Ancient dispersal of the human fungal pathogen *Cryptococcus gattii* from the Amazon rainforest. *PLoS ONE* **8**: e71148.
- Heckman DS, Geiser DM, Eidell BR, Stauffer RL, Kardos NL, Hedges SB. 2001. Molecular evidence for the early colonization of land by fungi and plants. *Science* **293**: 1129–1133.
- Hickman MA, Zeng G, Forche A, Hirakawa MP, Abbey D, Harrison BD, Wang YM, Su CH, Bennett RJ, Wang Y, et al. 2013. The “obligate diploid” *Candida albicans* forms mating-competent haploids. *Nature* **494**: 55–59.
- Hoff GL, Bigler WJ. 1981. The role of bats in the propagation and spread of histoplasmosis: A review. *J Wildl Dis* **17**: 191–196.
- Hoffmann K, Pawlowska J, Walther G, Wrzosek M, de Hoog GS, Benny GL, Kirk PM, Voigt K. 2013. The family structure of the *Mucorales*: A synoptic revision based on comprehensive multigene-genealogies. *Persoonia* **30**: 57–76.
- Hogan DA, Muhlschlegel FA. 2011. *Candida albicans* developmental regulation: Adenylyl cyclase as a coincidence detector of parallel signals. *Curr Opin Microbiol* **14**: 682–686.

J.R. Köhler et al.

- Holbrook ED, Rappleye CA. 2008. *Histoplasma capsulatum* pathogenesis: Making a lifestyle switch. *Curr Opin Microbiol* **11**: 318–324.
- Holland WL, Summers SA. 2008. Sphingolipids, insulin resistance, and metabolic disease: New insights from in vivo manipulation of sphingolipid metabolism. *Endocr Rev* **29**: 381–402.
- Hospenthal DR, Kwon-Chung KJ, Bennett JE. 1998. Concentrations of airborne *Aspergillus* compared to the incidence of invasive aspergillosis: Lack of correlation. *Med Mycol* **36**: 165–168.
- Hoyer LL, Green CB, Oh SH, Zhao X. 2008. Discovering the secrets of the *Candida albicans* agglutinin-like sequence (ALS) gene family—A sticky pursuit. *Med Mycol* **46**: 1–15.
- Hull CM, Johnson AD. 1999. Identification of a mating type-like locus in the asexual pathogenic yeast *Candida albicans*. *Science* **285**: 1271–1275.
- Humphreys CP, Franks PJ, Rees M, Bidartondo MI, Leake JR, Beerling DJ. 2010. Mutualistic mycorrhiza-like symbiosis in the most ancient group of land plants. *Nat Commun* **1**: 103.
- Ibrahim AS. 2011. Host cell invasion in mucormycosis: Role of iron. *Curr Opin Microbiol* **14**: 406–411.
- Ibrahim AS, Edwards JE Jr, Fu Y, Spellberg B. 2006. Deferiprone iron chelation as a novel therapy for experimental mucormycosis. *J Antimicrob Chemother* **58**: 1070–1073.
- Ibrahim AS, Spellberg B, Edwards J Jr. 2008. Iron acquisition: A novel perspective on mucormycosis pathogenesis and treatment. *Curr Opin Infect Dis* **21**: 620–625.
- Ibrahim AS, Gebremariam T, Lin L, Luo G, Hussein MI, Skory CD, Fu Y, French SW, Edwards JE Jr, Spellberg B. 2010. The high affinity iron permease is a key virulence factor required for *Rhizopus oryzae* pathogenesis. *Mol Microbiol* **77**: 587–604.
- Isa-Isa R, Arenas R, Fernandez RE, Isa M. 2012. Rhinofacial conidiobolomycosis (entomophthoromycosis). *Clin Dermatol* **30**: 409–412.
- Kam AP, Xu J. 2002. Diversity of commensal yeasts within and among healthy hosts. *Diagn Microbiol Infect Dis* **43**: 19–28.
- Kasuga T, Taylor JW, White TJ. 1999. Phylogenetic relationships of varieties and geographical groups of the human pathogenic fungus *Histoplasma capsulatum* Darling. *J Clin Microbiol* **37**: 653–663.
- Kasuga T, White TJ, Koenig G, McEwen J, Restrepo A, Castaneda E, Da Silva Lacaz C, Heins-Vaccari EM, De Freitas RS, Zancope-Oliveira RM, et al. 2003. Phylogeography of the fungal pathogen *Histoplasma capsulatum*. *Mol Ecol* **12**: 3383–3401.
- Kauffman CA. 2007. Histoplasmosis: A clinical and laboratory update. *Clin Microbiol Rev* **20**: 115–132.
- Kennedy AD, Willment JA, Dorward DW, Williams DL, Brown GD, DeLeo FR. 2007. Dectin-1 promotes fungicidal activity of human neutrophils. *Eur J Immunol* **37**: 467–478.
- Khan ZU, Ahmad S, Brazda A, Chandy R. 2009. *Mucor circinelloides* as a cause of invasive maxillofacial zygomycosis: An emerging dimorphic pathogen with reduced susceptibility to posaconazole. *J Clin Microbiol* **47**: 1244–1248.
- Kidd SE, Sorrell TC, Meyer W. 2003. Isolation of two molecular types of *Cryptococcus neoformans* var. *gattii* from insect frass. *Med Mycol* **41**: 171–176.
- Kidd SE, Hagen F, Tschärke RL, Huynh M, Bartlett KH, Fyfe M, Macdougall L, Boekhout T, Kwon-Chung KJ, Meyer W. 2004. A rare genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). *Proc Natl Acad Sci* **101**: 17258–17263.
- Kidd SE, Bach PJ, Hingston AO, Mak S, Chow Y, MacDougall L, Kronstad JW, Bartlett KH. 2007. *Cryptococcus gattii* dispersal mechanisms, British Columbia, Canada. *Emerg Infect Dis* **13**: 51–57.
- Kothary MH, Chase T Jr, Macmillan JD. 1984. Correlation of elastase production by some strains of *Aspergillus fumigatus* with ability to cause pulmonary invasive aspergillosis in mice. *Infect Immun* **43**: 320–325.
- Krcmery V, Barnes AJ. 2002. Non-albicans *Candida* spp. causing fungaemia: Pathogenicity and antifungal resistance. *J Hosp Infect* **50**: 243–260.
- Krockenberger MB, Canfield PJ, Malik R. 2002. *Cryptococcus neoformans* in the koala (*Phascolarctos cinereus*): Colonization by *C. n.* var. *gattii* and investigation of environmental sources. *Med Mycol* **40**: 263–272.
- Kronstad JW, Attarian R, Cadieux B, Choi J, D'Souza CA, Griffiths EJ, Geddes JM, Hu G, Jung WH, Kretschmer M, et al. 2011. Expanding fungal pathogenesis: *Cryptococcus* breaks out of the opportunistic box. *Nat Rev Microbiol* **9**: 193–203.
- Kronstad JW, Hu G, Jung WH. 2013. An encapsulation of iron homeostasis and virulence in *Cryptococcus neoformans*. *Trends Microbiol* **21**: 457–465.
- Kumamoto CA. 2008. Molecular mechanisms of mechanosensing and their roles in fungal contact sensing. *Nat Rev Microbiol* **6**: 667–673.
- Kwon-Chung KJ. 2012. Taxonomy of fungi causing mucormycosis and entomophthoromycosis (Zygomycosis) and nomenclature of the disease: Molecular mycologic perspectives. *Clin Infect Dis* **54**: S8–S15.
- Kwon-Chung KJ, Sugui JA. 2013. *Aspergillus fumigatus*—What makes the species a ubiquitous human fungal pathogen? *PLoS Pathog* **9**: e1003743.
- Latgé JP. 1999. *Aspergillus fumigatus* and aspergillosis. *Clin Microbiol Rev* **12**: 310–350.
- Lee SC, Li A, Calo S, Heitman J. 2013. Calcineurin plays key roles in the dimorphic transition and virulence of the human pathogenic zygomycete *Mucor circinelloides*. *PLoS Pathog* **9**: e1003625.
- Levitz SM, Diamond RD. 1985. Mechanisms of resistance of *Aspergillus fumigatus* Conidia to killing by neutrophils in vitro. *J Infect Dis* **152**: 33–42.
- Lin JS, Huang JH, Hung LY, Wu SY, Wu-Hsieh BA. 2010. Distinct roles of complement receptor 3, Dectin-1, and sialic acids in murine macrophage interaction with *Histoplasma* yeast. *J Leukoc Biol* **88**: 95–106.
- Litvintseva AP, Mitchell TG. 2012. Population genetic analyses reveal the African origin and strain variation of *Cryptococcus neoformans* var. *grubii*. *PLoS Pathog* **8**: e1002495.
- Liu Y, Filler SG. 2011. *Candida albicans* Als3, a multifunctional adhesin and invasin. *Eukaryot Cell* **10**: 168–173.

The Spectrum of Fungi That Infects Humans

- Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic genetic analysis of virulence in the human fungal pathogen *Cryptococcus neoformans*. *Cell* **135**: 174–188.
- Liu M, Spellberg B, Phan QT, Fu Y, Lee AS, Edwards JE Jr, Filler SG, Ibrahim AS. 2010. The endothelial cell receptor GRP78 is required for mucormycosis pathogenesis in diabetic mice. *J Clin Invest* **120**: 1914–1924.
- Lockhart SR, Wu W, Radke JB, Zhao R, Soll DR. 2005. Increased virulence and competitive advantage of a/a over a/a or α/α offspring conserves the mating system of *Candida albicans*. *Genetics* **169**: 1883–1890.
- Lott TJ, Fundyga RE, Kuykendall RJ, Arnold J. 2005. The human commensal yeast, *Candida albicans*, has an ancient origin. *Fungal Genet Biol* **42**: 444–451.
- Lucking R, Huhndorf S, Pfister DH, Plata ER, Lumsch HT. 2009. Fungi evolved right on track. *Mycologia* **101**: 810–822.
- Ma LJ, van der Does HC, Borkovich KA, Coleman JJ, Daboussi MJ, Di Pietro A, Dufresne M, Freitag M, Grabherr M, Henrissat B, et al. 2010. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* **464**: 367–373.
- Magee BB, Magee PT. 2000. Induction of mating in *Candida albicans* by construction of *MTLa* and *MTL α* strains. *Science* **289**: 310–313.
- Marples MJ, Di Menna ME. 1952. The incidence of *Candida albicans* in Dunedin, New Zealand. *J Pathol Bacteriol* **64**: 497–502.
- Marques SA. 2012. Paracoccidioidomycosis. *Clin Dermatol* **30**: 610–615.
- Marques SA. 2013. Paracoccidioidomycosis: Epidemiological, clinical, diagnostic and treatment up-dating. *An Bras Dermatol* **88**: 700–711.
- McCormick A, Heesemann L, Wagener J, Marcos V, Hartl D, Loeffler J, Heesemann J, Ebel F. 2010. NETs formed by human neutrophils inhibit growth of the pathogenic mold *Aspergillus fumigatus*. *Microbes Infect* **12**: 928–936.
- McGill S, Malik R, Saul N, Beetson S, Secombe C, Robertson I, Irwin P. 2009. Cryptococcosis in domestic animals in Western Australia: A retrospective study from 1995–2006. *Med Mycol* **47**: 625–639.
- McMurray DN, Greer DL. 1979. Immune responses in bats following intranasal infection with *Histoplasma capsulatum*. *Am J Trop Med Hyg* **28**: 1036–1039.
- Meyer W, Castaneda A, Jackson S, Huynh M, Castaneda E. 2003. Molecular typing of IberoAmerican *Cryptococcus neoformans* isolates. *Emerg Infect Dis* **9**: 189–195.
- Millanes AM, Truong C, Westberg M, Diederich P, Wedin M. 2014. Host switching promotes diversity in host-specialized mycoparasitic fungi: Uncoupled evolution in the *Biatoropsis-Usnea* system. *Evolution* doi:10.1111/evo.12374.
- Miller MG, Johnson AD. 2002. White-opaque switching in *Candida albicans* is controlled by mating-type locus homeodomain proteins and allows efficient mating. *Cell* **110**: 293–302.
- Miller WG, Padhye AA, van Bonn W, Jensen E, Brandt ME, Ridgway SH. 2002. Cryptococcosis in a bottlenose dolphin (*Tursiops truncatus*) caused by *Cryptococcus neoformans* var. *gattii*. *J Clin Microbiol* **40**: 721–724.
- Mitchell TG, Perfect JR. 1995. Cryptococcosis in the era of AIDS—100 years after the discovery of *Cryptococcus neoformans*. *Clin Microbiol Rev* **8**: 515–548.
- Molez JE. 1998. The historical question of acquired immunodeficiency syndrome in the 1960s in the Congo River basin area in relation to cryptococcal meningitis. *Am J Trop Med Hyg* **58**: 273–276.
- Money NB, Davis CM, Ravishanker JP. 2004. Biomechanical evidence for convergent evolution of the invasive growth process among fungi and oomycete water molds. *Fungal Genet Biol* **41**: 872–876.
- Morita T, Kishimoto M, Shimada A, Matsumoto Y, Shindo J. 2001. Disseminated histoplasmosis in a sea otter (*Enhydra lutris*). *J Comp Pathol* **125**: 219–223.
- Morschhäuser J. 2011. Nitrogen regulation of morphogenesis and protease secretion in *Candida albicans*. *Int J Med Microbiol* **301**: 390–394.
- Morton CO, Bouzani M, Loeffler J, Rogers TR. 2012. Direct interaction studies between *Aspergillus fumigatus* and human immune cells: What have we learned about pathogenicity and host immunity? *Front Microbiol* **3**: 413.
- Mullins J, Harvey R, Seaton A. 1976. Sources and incidence of airborne *Aspergillus fumigatus* (Fres). *Clin Allergy* **6**: 209–217.
- Neilson JB, Ivey MH, Bulmer GS. 1978. *Cryptococcus neoformans*: Pseudohyphal forms surviving culture with *Acanthamoeba polyphaga*. *Infect Immun* **20**: 262–266.
- Nemecek JC, Wuthrich M, Klein BS. 2006. Global control of dimorphism and virulence in fungi. *Science* **312**: 583–588.
- Netea MG, Warris A, Van der Meer JW, Fenton MJ, Verver-Janssen TJ, Jacobs LE, Andresen T, Verweij PE, Kullberg BJ. 2003. *Aspergillus fumigatus* evades immune recognition during germination through loss of toll-like receptor-4-mediated signal transduction. *J Infect Dis* **188**: 320–326.
- Nguyen VQ, Sil A. 2008. Temperature-induced switch to the pathogenic yeast form of *Histoplasma capsulatum* requires Ryp1, a conserved transcriptional regulator. *Proc Natl Acad Sci* **105**: 4880–4885.
- Nguyen C, Barker BM, Hoover S, Nix DE, Ampel NM, Frelinger JA, Orbach MJ, Galgiani JN. 2013. Recent advances in our understanding of the environmental, epidemiological, immunological, and clinical dimensions of coccidioidomycosis. *Clin Microbiol Rev* **26**: 505–525.
- Nicol MR, Adams JL, Kashuba AD. 2013. HIV PrEP trials: The road to success. *Clin Invest (Lond)* **3**: 295–308.
- Nielsen K, De Obaldia AL, Heitman J. 2007. *Cryptococcus neoformans* mates on pigeon guano: Implications for the realized ecological niche and globalization. *Eukaryot Cell* **6**: 949–959.
- Noble SM. 2013. *Candida albicans* specializations for iron homeostasis: From commensalism to virulence. *Curr Opin Microbiol* **16**: 708–715.
- Noble SM, French S, Kohn LA, Chen V, Johnson AD. 2010. Systematic screens of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and pathogenicity. *Nat Genet* **42**: 590–598.
- Nucci M, Anaissie E. 2007. *Fusarium* infections in immunocompromised patients. *Clin Microbiol Rev* **20**: 695–704.

J.R. Köhler et al.

- O'Brien HE, Parrent JL, Jackson JA, Moncalvo JM, Vilgalys R. 2005. Fungal community analysis by large-scale sequencing of environmental samples. *Appl Environ Microbiol* **71**: 5544–5550.
- Odds FC. 1984. Ecology and epidemiology of *Candida* species. *Zentralbl Bakteriol Mikrobiol Hyg A* **257**: 207–212.
- Odds F. 1988. *Candida and candidosis: A review and bibliography*. Bailliere and Tindall, London.
- O'Donnell K, Sutton DA, Rinaldi MG, Magnon KC, Cox PA, Revankar SG, Sanche S, Geiser DM, Juba JH, van Burik JA, et al. 2004. Genetic diversity of human pathogenic members of the *Fusarium oxysporum* complex inferred from multilocus DNA sequence data and amplified fragment length polymorphism analyses: Evidence for the recent dispersion of a geographically widespread clonal lineage and nosocomial origin. *J Clin Microbiol* **42**: 5109–5120.
- Okafor JI. 1994. Purification and characterization of protease enzymes of *Basidiobolus* and *Conidiobolus* species. *Mycoses* **37**: 265–269.
- Okafor JI, Gugnani HC. 1990. Lipase activity of *Basidiobolus* and *Conidiobolus* species. *Mycoses* **33**: 81–85.
- Okafor JI, Gugnani HC, Testratke D, Yangoo BG. 1987. Extracellular enzyme activities by *Basidiobolus* and *Conidiobolus* isolates on solid media. *Mykosen* **30**: 404–407.
- Okagaki LH, Strain AK, Nielsen JN, Charlier C, Baltes NJ, Chretien F, Heitman J, Dromer F, Nielsen K. 2010. Cryptococcal cell morphology affects host cell interactions and pathogenicity. *PLoS Pathog* **6**: e1000953.
- Oliveira DL, Nakayasu ES, Joffe LS, Guimaraes AJ, Sobreira TJ, Nosanchuk JD, Cordero RJ, Frases S, Casadevall A, Almeida IC, et al. 2010. Characterization of yeast extracellular vesicles: Evidence for the participation of different pathways of cellular traffic in vesicle biogenesis. *PLoS ONE* **5**: e11113.
- Paris S, Debeaupuis JP, Cramer R, Carey M, Charles F, Prevost MC, Schmitt C, Philippe B, Latgé JP. 2003. Conidial hydrophobins of *Aspergillus fumigatus*. *Appl Environ Microbiol* **69**: 1581–1588.
- Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM. 2009. Estimation of the current global burden of cryptococcal meningitis among persons living with HIV/AIDS. *AIDS* **23**: 525–530.
- Perera TH, Gregory DW, Marshall D, Gow NA. 1997. Contact-sensing by hyphae of dermatophytic and saprophytic fungi. *J Med Vet Mycol* **35**: 289–293.
- Pfaller MA, Diekema DJ. 2007. Epidemiology of invasive candidiasis: A persistent public health problem. *Clin Microbiol Rev* **20**: 133–163.
- Pihet M, Vandeputte P, Tronchin G, Renier G, Saulnier P, Georgeault S, Mallet R, Chabasse D, Symoens F, Bouchara JP. 2009. Melanin is an essential component for the integrity of the cell wall of *Aspergillus fumigatus* conidia. *BMC Microbiol* **9**: 177.
- Poole CJ, Wong M. 2013. Allergic bronchopulmonary aspergillosis in garden waste (compost) collectors—Occupational implications. *Occup Med (Lond)* **63**: 517–519.
- Prabhu RM, Patel R. 2004. Mucormycosis and entomophthoromycosis: A review of the clinical manifestations, diagnosis and treatment. *Clin Microbiol Infect* **10**: 31–47.
- Puccia R, Juliano MA, Juliano L, Travassos LR, Carmona AK. 1999. Detection of the basement membrane-degrading proteolytic activity of *Paracoccidioides brasiliensis* after SDS-PAGE using agarose overlays containing Abz-MKALTQ-EDDnp. *Braz J Med Biol Res* **32**: 645–649.
- Puccia R, McEwen JG, Cicalino PS. 2008. Diversity in *Paracoccidioides brasiliensis*. The *PbGP43* gene as a genetic marker. *Mycopathologia* **165**: 275–287.
- Puccia R, Vallejo MC, Matsuo AL, Longo IV. 2011. The *paracoccidioides* cell wall: Past and present layers toward understanding interaction with the host. *Front Microbiol* **2**: 257.
- Pukkila-Worley R, Mylonakis E. 2008. Epidemiology and management of cryptococcal meningitis: Developments and challenges. *Expert Opin Pharmacother* **9**: 551–560.
- Quan C, Spellberg B. 2010. Mucormycosis, pseudallescheriasis, and other uncommon mold infections. *Proc Am Thorac Soc* **7**: 210–215.
- Queiroz-Telles F, Escuissato DL. 2011. Pulmonary paracoccidioidomycosis. *Semin Respir Crit Care Med* **32**: 764–774.
- Quist EM, Belcher C, Levine G, Johnson M, Heatley JJ, Kiupel M, Giri D. 2011. Disseminated histoplasmosis with concurrent oral candidiasis in an Eclectus parrot (*Eclectus roratus*). *Avian Pathol* **40**: 207–211.
- Rappleye CA, Engle JT, Goldman WE. 2004. RNA interference in *Histoplasma capsulatum* demonstrates a role for $\alpha(1,3)$ -glucan in virulence. *Mol Microbiol* **53**: 153–165.
- Rappleye CA, Eissenberg LG, Goldman WE. 2007. *Histoplasma capsulatum* $\alpha(1,3)$ -glucan blocks innate immune recognition by the β -glucan receptor. *Proc Natl Acad Sci* **104**: 1366–1370.
- Raso TF, Werther K, Miranda ET, Mendes-Giannini MJ. 2004. Cryptococcosis outbreak in psittacine birds in Brazil. *Med Mycol* **42**: 355–362.
- Redecker D, Kodner R, Graham LE. 2000. Glomalean fungi from the Ordovician. *Science* **289**: 1920–1921.
- Rep M, Kistler HC. 2010. The genomic organization of plant pathogenicity in *Fusarium* species. *Curr Opin Plant Biol* **13**: 420–426.
- Ribes JA, Vanover-Sams CL, Baker DJ. 2000. Zygomycetes in human disease. *Clin Microbiol Rev* **13**: 236–301.
- Richini-Pereira VB, Bosco Sde M, Theodoro RC, Macoris SA, Bagagli E. 2009. Molecular approaches for eco-epidemiological studies of *Paracoccidioides brasiliensis*. *Mem Inst Oswaldo Cruz* **104**: 636–643.
- Robert VA, Casadevall A. 2009. Vertebrate endothermy restricts most fungi as potential pathogens. *J Infect Dis* **200**: 1623–1626.
- Roden MM, Zaoutis TE, Buchanan WL, Knudsen TA, Sarkisova TA, Schaufele RL, Sein M, Sein T, Chiou CC, Chu JH, et al. 2005. Epidemiology and outcome of zygomycosis: A review of 929 reported cases. *Clin Infect Dis* **41**: 634–653.
- Rodrigues ML, Nimrichter L, Oliveira DL, Frases S, Miranda K, Zaragoza O, Alvarez M, Nakouzi A, Feldmesser M, Casadevall A. 2007. Vesicular polysaccharide export in *Cryptococcus neoformans* is a eukaryotic solution to the problem of fungal trans-cell wall transport. *Eukaryot Cell* **6**: 48–59.

- Rodrigues ML, Nakayasu ES, Oliveira DL, Nimrichter L, Nosanchuk JD, Almeida IC, Casadevall A. 2008. Extracellular vesicles produced by *Cryptococcus neoformans* contain protein components associated with virulence. *Eukaryot Cell* **7**: 58–67.
- Rohde JR, Cardenas ME. 2004. Nutrient signaling through TOR kinases controls gene expression and cellular differentiation in fungi. *Curr Top Microbiol Immunol* **279**: 53–72.
- Rohm M, Grimm MJ, D'Auria AC, Almyroudis NG, Segal BH, Urban CF. 2014. NADPH oxidase promotes neutrophil extracellular trap formation in pulmonary aspergillosis. *Infect Immun* **82**: 1766–1777.
- Saijo T, Chen J, Chen SC, Rosen LB, Yi J, Sorrell TC, Bennett JE, Holland SM, Browne SK, Kwon-Chung KJ. 2014. Anti-granulocyte-macrophage colony-stimulating factor autoantibodies are a risk factor for central nervous system infection by *Cryptococcus gattii* in otherwise immunocompetent patients. *mBio* **5**: e00912-14.
- Santos WR, Meyer W, Wanke B, Costa SP, Trilles L, Nascimento JL, Medeiros R, Morales BP, Bezerra Cde C, Macedo RC, et al. 2008. Primary endemic *Cryptococcosis gattii* by molecular type VGII in the state of Para, Brazil. *Mem Inst Oswaldo Cruz* **103**: 813–818.
- Schaller M, Borelli C, Korting HC, Hube B. 2005. Hydrolytic enzymes as virulence factors of *Candida albicans*. *Mycoses* **48**: 365–377.
- Schmitt HJ, Blevins A, Soback K, Armstrong D. 1990. *Aspergillus* species from hospital air and from patients. *Mycoses* **33**: 539–541.
- Sebghati TS, Engle JT, Goldman WE. 2000. Intracellular parasitism by *Histoplasma capsulatum*: Fungal virulence and calcium dependence. *Science* **290**: 1368–1372.
- Shankar J, Restrepo A, Clemons KV, Stevens DA. 2011. Hormones and the resistance of women to paracoccidioidomycosis. *Clin Microbiol Rev* **24**: 296–313.
- Shapiro RS, Ryan O, Boone C, Cowen LE. 2012. Regulatory circuitry governing morphogenesis in *Saccharomyces cerevisiae* and *Candida albicans*. *Cell Cycle* **11**: 4294–4295.
- Shen J, Cowen LE, Griffin AM, Chan L, Köhler JR. 2008. The *Candida albicans* *pescadillo* homolog is required for normal hypha-to-yeast morphogenesis and yeast proliferation. *Proc Natl Acad Sci* **105**: 20918–20923.
- Shi M, Li SS, Zheng C, Jones GJ, Kim KS, Zhou H, Kubes P, Mody CH. 2010. Real-time imaging of trapping and urease-dependent transmigration of *Cryptococcus neoformans* in mouse brain. *J Clin Invest* **120**: 1683–1693.
- Shiraki Y, Ishibashi Y, Hiruma M, Nishikawa A, Ikeda S. 2006. Cytokine secretion profiles of human keratinocytes during *Trichophyton tonsurans* and *Arthroderma benhamiae* infections. *J Med Microbiol* **55**: 1175–1185.
- Siegfried N, van der Merwe L, Brocklehurst P, Sint TT. 2011. Antiretrovirals for reducing the risk of mother-to-child transmission of HIV infection. *Cochrane Database Syst Rev* CD003510.
- Simwami SP, Khayhan K, Henk DA, Aanensen DM, Boekhout T, Hagen F, Brouwer AE, Harrison TS, Donnelly CA, Fisher MC. 2011. Low diversity *Cryptococcus neoformans* variety *grubii* multilocus sequence types from Thailand are consistent with an ancestral African origin. *PLoS Pathog* **7**: e1001343.
- Skinner CE, Fletcher DW. 1960. A review of the genus *Candida*. *Bacteriol Rev* **24**: 397–416.
- Smith RM, Schaefer MK, Kainer MA, Wise M, Finks J, Duwve J, Fontaine E, Chu A, Carothers B, Reilly A, et al. 2013. Fungal infections associated with contaminated methylprednisolone injections. *N Engl J Med* **369**: 1598–1609.
- Soll DR. 2002. *Candida* commensalism and virulence: The evolution of phenotypic plasticity. *Acta Trop* **81**: 101–110.
- Soll DR, Galask R, Schmid J, Hanna C, Mac K, Morrow B. 1991. Genetic dissimilarity of commensal strains of *Candida* spp. carried in different anatomical locations of the same healthy women. *J Clin Microbiol* **29**: 1702–1710.
- Speed B, Dunt D. 1995. Clinical and host differences between infections with the two varieties of *Cryptococcus neoformans*. *Clin Infect Dis* **21**: 28–34; discussion 35–36.
- Spellberg B, Edwards J Jr, Ibrahim A. 2005. Novel perspectives on mucormycosis: Pathophysiology, presentation, and management. *Clin Microbiol Rev* **18**: 556–569.
- Spellberg B, Andes D, Perez M, Anglim A, Bonilla H, Mathisen GE, Walsh TJ, Ibrahim AS. 2009. Safety and outcomes of open-label deferasirox iron chelation therapy for mucormycosis. *Antimicrob Agents Chemother* **53**: 3122–3125.
- Springer DJ, Phadke S, Billmyre B, Heitman J. 2012. *Cryptococcus gattii*, no longer an accidental pathogen? *Curr Fungal Infect Rep* **6**: 245–256.
- Staab JE, Bradway SD, Fidel PL, Sundstrom P. 1999. Adhesive and mammalian transglutaminase substrate properties of *Candida albicans* Hwp1. *Science* **283**: 1535–1538.
- Staib P, Kretschmar M, Nichterlein T, Hof H, Morschhäuser J. 2000. Differential activation of a *Candida albicans* virulence gene family during infection. *Proc Natl Acad Sci* **97**: 6102–6107.
- Steenbergen JN, Shuman HA, Casadevall A. 2001. *Cryptococcus neoformans* interactions with amoebae suggest an explanation for its virulence and intracellular pathogenic strategy in macrophages. *Proc Natl Acad Sci* **98**: 15245–15250.
- Steenbergen JN, Nosanchuk JD, Malliaris SD, Casadevall A. 2003. *Cryptococcus neoformans* virulence is enhanced after growth in the genetically malleable host *Dictyostelium discoideum*. *Infect Immun* **71**: 4862–4872.
- Steenbergen JN, Nosanchuk JD, Malliaris SD, Casadevall A. 2004. Interaction of *Blastomyces dermatitidis*, *Sporothrix schenckii*, and *Histoplasma capsulatum* with *Acanthamoeba castellanii*. *Infect Immun* **72**: 3478–3488.
- Sudbery P, Gow N, Berman J. 2004. The distinct morphogenic states of *Candida albicans*. *Trends Microbiol* **12**: 317–324.
- Sundstrom P. 2002. Adhesion in *Candida* spp. *Cell Microbiol* **4**: 461–469.
- Taylor ML, Chavez-Tapia CB, Vargas-Yanez R, Rodriguez-Arellanes G, Pena-Sandoval GR, Toriello C, Perez A, Reyes-Montes MR. 1999. Environmental conditions favoring bat infection with *Histoplasma capsulatum* in Mexican shelters. *Am J Trop Med Hyg* **61**: 914–919.
- Thau N, Monod M, Crestani B, Rolland C, Tronchin G, Latgé JP, Paris S. 1994. *rodletless* mutants of *Aspergillus fumigatus*. *Infect Immun* **62**: 4380–4388.

J.R. Köhler et al.

- Theodoro RC, Teixeira Mde M, Felipe MS, Paduan Kdos S, Ribolla PM, San-Blas G, Bagagli E. 2012. Genus *paracoccidioides*: Species recognition and biogeographic aspects. *PLoS ONE* **7**: e37694.
- Torres HA, Rivero GA, Lewis RE, Hachem R, Raad II, Kontoyiannis DP. 2003. Aspergillosis caused by non-*fumigatus* *Aspergillus* species: Risk factors and in vitro susceptibility compared with *Aspergillus fumigatus*. *Diagn Microbiol Infect Dis* **46**: 25–28.
- Trofa D, Soghier L, Long C, Nosanchuk JD, Gacser A, Goldman DL. 2011. A rat model of neonatal candidiasis demonstrates the importance of lipases as virulence factors for *Candida albicans* and *Candida parapsilosis*. *Mycopathologia* **172**: 169–178.
- Tucker SC, Casadevall A. 2002. Replication of *Cryptococcus neoformans* in macrophages is accompanied by phagosomal permeabilization and accumulation of vesicles containing polysaccharide in the cytoplasm. *Proc Natl Acad Sci* **99**: 3165–3170.
- Underhill DM, Rossnagle E, Lowell CA, Simmons RM. 2005. Dectin-1 activates Syk tyrosine kinase in a dynamic subset of macrophages for reactive oxygen production. *Blood* **106**: 2543–2550.
- Untereiner WA, Scott JA, Naveau FA, Sigler L, Bachewich J, Angus A. 2004. The Ajellomycetaceae, a new family of vertebrate-associated Onygenales. *Mycologia* **96**: 812–821.
- Vallejo MC, Matsuo AL, Ganiko L, Medeiros LC, Miranda K, Silva LS, Freymuller-Haapalainen E, Sinigaglia-Coimbra R, Almeida IC, Puccia R. 2011. The pathogenic fungus *Paracoccidioides brasiliensis* exports extracellular vesicles containing highly immunogenic α -galactosyl epitopes. *Eukaryot Cell* **10**: 343–351.
- Vallejo MC, Nakayasu ES, Matsuo AL, Sobreira TJ, Longo IV, Ganiko L, Almeida IC, Puccia R. 2012. Vesicle and vesicle-free extracellular proteome of *Paracoccidioides brasiliensis*: Comparative analysis with other pathogenic fungi. *J Proteome Res* **11**: 1676–1685.
- Vikram HR, Smilack JD, Leighton JA, Crowell MD, De Petris G. 2012. Emergence of gastrointestinal basidiobolomycosis in the United States, with a review of worldwide cases. *Clin Infect Dis* **54**: 1685–1691.
- Vilela R, Rosa PS, Belone AF, Taylor JW, Diorio SM, Mendoza L. 2009. Molecular phylogeny of animal pathogen *Lacazia loboi* inferred from rDNA and DNA coding sequences. *Mycol Res* **113**: 851–857.
- Wang Y, Aisen P, Casadevall A. 1995. *Cryptococcus neoformans* melanin and virulence: Mechanism of action. *Infect Immun* **63**: 3131–3136.
- Wasylnka JA, Moore MM. 2003. *Aspergillus fumigatus* conidia survive and germinate in acidic organelles of A549 epithelial cells. *J Cell Sci* **116**: 1579–1587.
- Webster RH, Sil A. 2008. Conserved factors Ryp2 and Ryp3 control cell morphology and infectious spore formation in the fungal pathogen *Histoplasma capsulatum*. *Proc Natl Acad Sci* **105**: 14573–14578.
- Weissman Z, Shemer R, Conibear E, Kornitzer D. 2008. An endocytic mechanism for haemoglobin-iron acquisition in *Candida albicans*. *Mol Microbiol* **69**: 201–217.
- Weitzman I, Summerbell RC. 1995. The dermatophytes. *Clin Microbiol Rev* **8**: 240–259.
- Werner JL, Metz AE, Horn D, Schoeb TR, Hewitt MM, Schwiebert LM, Faro-Trindade I, Brown GD, Steele C. 2009. Requisite role for the dectin-1 β -glucan receptor in pulmonary defense against *Aspergillus fumigatus*. *J Immunol* **182**: 4938–4946.
- Wheeler RT, Fink GR. 2006. A drug-sensitive genetic network masks fungi from the immune system. *PLoS Pathog* **2**: e35.
- Wieloch W, Bogus MI, Ligeza M, Koszela-Piotrowska I, Szewczyk A. 2011. Coronatin-1 isolated from entomopathogenic fungus *Conidiobolus coronatus* kills *Galleria mellonella* hemocytes in vitro and forms potassium channels in planar lipid membrane. *Toxicon* **58**: 369–379.
- Williams MJ. 2007. Drosophila hemopoiesis and cellular immunity. *J Immunol* **178**: 4711–4716.
- Windus DW, Stokes TJ, Julian BA, Fenves AZ. 1987. Fatal Rhizopus infections in hemodialysis patients receiving deferroxamine. *Ann Intern Med* **107**: 678–680.
- Wrobel L, Whittington JK, Pujol C, Oh SH, Ruiz MO, Pfaller MA, Diekema DJ, Soll DR, Hoyer LL. 2008. Molecular phylogenetic analysis of a geographically and temporally matched set of *Candida albicans* isolates from humans and nonmigratory wildlife in central Illinois. *Eukaryot Cell* **7**: 1475–1486.
- Wu W, Lockhart SR, Pujol C, Srikantha T, Soll DR. 2007. Heterozygosity of genes on the sex chromosome regulates *Candida albicans* virulence. *Mol Microbiol* **64**: 1587–1604.
- Wu Y, Yang J, Yang F, Liu T, Leng W, Chu Y, Jin Q. 2009. Recent dermatophyte divergence revealed by comparative and phylogenetic analysis of mitochondrial genomes. *BMC Genomics* **10**: 238.
- Youseff BH, Dougherty JA, Rappleye CA. 2009. Reverse genetics through random mutagenesis in *Histoplasma capsulatum*. *BMC Microbiol* **9**: 236.
- Zaragoza O, Garcia-Rodas R, Nosanchuk JD, Cuenca-Estrella M, Rodriguez-Tudela JL, Casadevall A. 2010. Fungal cell gigantism during mammalian infection. *PLoS Pathog* **6**: e1000945.
- Zavasky DM, Samowitz W, Loftus T, Segal H, Carroll K. 1999. Gastrointestinal zygomycotic infection caused by *Basidiobolus ranarum*: Case report and review. *Clin Infect Dis* **28**: 1244–1248.
- Zimmerman LE. 1955. Fatal fungus infections complicating other diseases. *Am J Clin Pathol* **25**: 46–65.
- Zugmaier W, Bauer R, Oberwinkler F. 1994. Mycoparasitism of some *Tremella* species. *Mycologia* **86**: 49–56.

The Spectrum of Fungi That Infects Humans

Julia R. Köhler, Arturo Casadevall and John Perfect

Cold Spring Harb Perspect Med 2015; doi: 10.1101/cshperspect.a019273 originally published online November 3, 2014

Subject Collection [Human Fungal Pathogens](#)

Evolutionary Perspectives on Human Fungal Pathogens

John W. Taylor

Black Molds and Melanized Yeasts Pathogenic to Humans

Anuradha Chowdhary, John Perfect and G. Sybren de Hoog

Fungal Pathogens: Survival and Replication within Macrophages

Andrew S. Gilbert, Robert T. Wheeler and Robin C. May

Innate Defense against Fungal Pathogens

Rebecca A. Drummond, Sarah L. Gaffen, Amy G. Hise, et al.

Antifungal Pharmacokinetics and Pharmacodynamics

Alexander J. Lepak and David R. Andes

Human Fungal Pathogens of Mucorales and Entomophthorales

Leonel Mendoza, Raquel Vilela, Kerstin Voelz, et al.

Functional Profiling of Human Fungal Pathogen Genomes

Alexi I. Goranov and Hiten D. Madhani

***Aspergillus fumigatus* and Related Species**

Janyce A. Sugui, Kyung J. Kwon-Chung, Praveen R. Juvvadi, et al.

Thermally Dimorphic Human Fungal Pathogens—Polyphyletic Pathogens with a Convergent Pathogenicity Trait

Anita Sil and Alex Andrianopoulos

Mechanisms of Antifungal Drug Resistance

Leah E. Cowen, Dominique Sanglard, Susan J. Howard, et al.

Treatment Principles for *Candida* and *Cryptococcus*

Laura C. Whitney and Tihana Bicanic

The Human Mycobiome

Patrick C. Seed

Treatment Principles for the Management of Mold Infections

Dimitrios P. Kontoyiannis and Russell E. Lewis

Adaptive Immunity to Fungi

Akash Verma, Marcel Wüthrich, George Deepe, et al.

The *Candida* Pathogenic Species Complex

Siobhán A. Turner and Geraldine Butler

Fungal Morphogenesis

Xiaorong Lin, J. Andrew Alspaugh, Haoping Liu, et al.

For additional articles in this collection, see <http://perspectivesinmedicine.cshlp.org/cgi/collection/>